

NON-DISCRETE EUCLIDEAN BUILDINGS FOR THE REE AND SUZUKI GROUPS

PETRA HITZELBERGER, LINUS KRAMER AND RICHARD M. WEISS

ABSTRACT. We call a non-discrete Euclidean building a *Bruhat-Tits space* if its automorphism group contains a subgroup that induces the subgroup generated by all the root groups of a root datum of the building at infinity. This is the class of non-discrete Euclidean buildings introduced and studied by Bruhat and Tits in [2]. We give the complete classification of Bruhat-Tits spaces whose building at infinity is the fixed point set of a polarity of an ambient building of type B_2 , F_4 or G_2 associated with a Ree or Suzuki group endowed with the usual root datum. (In the B_2 and G_2 cases, this fixed point set is a building of rank one; in the F_4 case, it is a generalized octagon whose Weyl group is not crystallographic.) We also show that each of these Bruhat-Tits spaces has a natural embedding in the unique Bruhat-Tits space whose building at infinity is the corresponding ambient building.

1. INTRODUCTION

Suppose that X is an irreducible affine building. Typically (and for certain if the dimension of X is at least three), the automorphism group of the building at infinity of X contains subgroups constituting a *root datum* defined over a field K . In this case the affine building X is uniquely determined by a valuation of this root datum which is, in turn, uniquely determined by a discrete valuation of K .

The notion of a valuation of the root datum of a spherical building makes perfectly good sense if we drop the requirement that the values lie in a discrete subgroup of $\mathbb{R}$. In the non-discrete case, there is no longer a corresponding affine building X . There does exist, however, an analogous structure called a *non-discrete Euclidean building*.

Non-discrete Euclidean buildings were first introduced and studied by Bruhat and Tits in [2]. These structures were first axiomatized and, in dimension greater than two, classified by Tits in [12]. (Other fundamental references about non-discrete Euclidean buildings are [4] and [6]; see also [1] and [8].)

Non-discrete Euclidean buildings are sometimes called *affine $\mathbb{R}$ -buildings* (since in the simplest case they are $\mathbb{R}$ -trees) or *apartment systems* (since they have apartments but are not really buildings) or, as in [4], simply *Euclidean buildings*, although this term is more commonly synonymous with “affine building.”

In 4.13 below, we propose the term *Bruhat-Tits space* for the class of non-discrete Euclidean buildings that were introduced and studied by Bruhat and Tits in [2]. (The term *Bruhat-Tits space* was used in [5] to describe complete metric spaces that

Address of the first two authors: Fachbereich Mathematik und Informatik, Universität Münster, Einsteinstrasse 62, 48149 Münster, Germany.

Address of the third author: Department of Mathematics, Tufts University, 503 Boston Avenue, Medford, MA 02155, USA.

satisfy a certain “semi-parallelogram rule” introduced by S. Lang. This is a more general class of metric spaces which includes not only all non-discrete Euclidean buildings but also, for example, all simply connected Riemannian manifolds of non-positive curvature. We mention too that Bruhat-Tits spaces in our sense are not necessarily complete as metric spaces; see Section 7.5 in [2].)

Suppose now that Δ is the spherical building (of rank one or two) associated with a Ree or Suzuki group. Thus Δ is the fixed point set of a “polarity” of a building of type B_2 , F_4 or G_2 which is defined over a field extension K/F , where $p := \text{char}(K)$ equals 2 in the first two cases and 3 in the third case, and the polarity is defined in terms of a Tits endomorphism θ of K (as defined in 5.1) whose image is F . On page 173 of [12], Tits remarks that an arbitrary valuation ν of K (where “arbitrary” means “arbitrary non-trivial real-valued non-archimedean”) extends to a valuation of the root datum of Δ —and thus there exists a corresponding Bruhat-Tits space whose building at infinity is Δ —if and only if the valuation ν is θ -invariant. It is the goal of this paper to make this statement more precise and to fill in all the details justifying it.

Here θ -invariant means that $\nu \circ \theta$ is equivalent to ν , i.e. that $\nu(x) \geq 0$ for $x \in K^*$ if and only if $\nu(x^\theta) \geq 0$. This is the same as saying that $\nu(x^\theta) = \gamma \cdot \nu(x)$ for some positive real number γ and all $x \in K^*$. Since θ is a Tits endomorphism of K , it follows that ν is θ -invariant if and only if

$$\nu(x^\theta) = \sqrt{p} \cdot \nu(x)$$

for all $x \in K^*$. Thus, in particular, ν cannot be θ -invariant if it is discrete (since the ratio of two values of a discrete valuation is always rational).

2. OVERVIEW

Continuing to put precision aside for a moment, we can summarize the main results of this paper roughly as follows; see also 8.11.

Theorem 2.1. *Let G be a Ree or Suzuki group. Then the following hold:*

- (i) *There exists a Moufang building $\dot{\Delta}$ of type B_2 , F_4 or G_2 having a polarity ρ defined over a pair (K, θ) as described in Section 1 such that G is the group induced by the centralizer $C_{\dot{G}}(\rho)$ on the set $\dot{\Delta}^\rho$ of fixed points of ρ , where $\dot{G}$ is the subgroup of $\text{Aut}(\dot{\Delta})$ generated by the root groups of $\dot{\Delta}$.*
- (ii) *The set $\Delta := \dot{\Delta}^\rho$ has the structure of a Moufang building of rank one in cases B_2 and G_2 , of rank two whose Weyl group is dihedral of order 16 in case F_4 .*
- (iii) *For each valuation ν of the field K , there exists a unique non-discrete Euclidean building $(\dot{X}, \dot{A})$ determined by ν whose building at infinity is $\dot{\Delta}$ and whose automorphism group induces $\dot{G}$ on $\dot{\Delta}$.*
- (iv) *Let ν and $(\dot{X}, \dot{A})$ be as in (iii). Then there exists an automorphism $\dot{\rho}$ of $(\dot{X}, \dot{A})$ inducing the polarity ρ on $\dot{\Delta}$ if and only if ν is θ -invariant. Furthermore, $\dot{\rho}$, if it exists, is unique.*
- (v) *If ν , $(\dot{X}, \dot{A})$ and $\dot{\rho}$ are as in (iv), then there is a canonical non-discrete Euclidean building (X, A) contained in the fixed point set of $\dot{\rho}$ in $(\dot{X}, \dot{A})$ whose building at infinity is Δ and whose automorphism group contains a subgroup inducing G on Δ .*

- (vi) *Every non-discrete Euclidean building $(X, \mathcal{A})$ whose building at infinity is Δ and whose automorphism group contains a subgroup inducing G on Δ arises from a θ -invariant valuation ν of K as described in (v).*

Proof. Assertion (i) is essentially the definition of a Ree or Suzuki group. The building $\dot{\Delta}$ and its polarity ρ are described in 5.3. Assertion (ii) is proved in 6.5.iii and 6.7. Assertions (iii) and (iv) follow from 4.16 and 5.15. Assertion (v) is proved in 7.11 and assertion (vi) is a consequence of 7.1. $\square$

This paper is organized as follows: In Sections 3–4 we review the basic results about root data, valuations of root data and non-discrete Euclidean buildings we require. In Sections 5 we introduce the spherical buildings of type B_2 , F_4 and G_2 having polarities that give rise to the Ree and Suzuki groups and in Section 6 we assemble the basic properties of the subbuildings fixed by these polarities we require. Our main results—7.1, 7.4 and 7.11—are then proved in Sections 7 and 8.

Throughout this paper we will use Δ and similar letters to denote spherical buildings and $(X, \mathcal{A})$ and similar letters to denote Euclidean buildings.

3. ROOT DATA AND VALUATIONS

We now start paying attention to the details. In this section we review the notions of a root datum of a spherical building and a valuation of a root datum.

Notation 3.1. Let (W, S) be an irreducible spherical Coxeter system. Then either W can be identified with the Weyl group of an irreducible root system Φ so that S consists of the reflections determined by the elements in a basis, or $|S| = 2$ and W is a dihedral group of order $2n$ for $n = 5$ or $n > 6$. In the latter case we let Φ consist of $2n$ vectors evenly distributed around the unit circle in a 2-dimensional Euclidean space and think of S as the two reflections determined by two of these vectors making an angle of $(n - 1)180/n$ degrees. (Later we will refer to this set Φ as $I_2(n)$.) In both cases, we denote by $\mathbb{A}$ the ambient Euclidean space of Φ and by $\text{Aut}(\Phi)$ the group of isometries of $\mathbb{A}$ mapping Φ to itself. Note that (W, S) is uniquely determined by Φ . When we sometimes call W the *Weyl group* of Φ (as is usual), we really have the pair (W, S) in mind.

Notation 3.2. Let (W, S) and Φ be as in 3.1. For all $\alpha, \beta \in \Phi$ such that $\alpha \neq \pm\beta$, the *interval* (α, β) is the s -tuple $(\gamma_1, \gamma_2, \dots, \gamma_s)$ of all elements $\gamma_i \in \Phi$ such that for some positive real numbers p_i and q_i (which depend on α and β),

$$(3.3) \quad \gamma_i / |\gamma_i| = p_i \alpha / |\alpha| + q_i \beta / |\beta|,$$

where $\angle(\alpha, \gamma_i) < \angle(\alpha, \gamma_j)$ if and only if $i < j$. Note that s depends on the pair α and β and that for some pairs, $s = 0$.

To define the interval (α, β) , we could have omitted the denominators in 3.3. We included the denominators because the coefficients p_i and q_i in this equation (as it is written) are needed in the statement of condition (V2) in 3.14.

Notation 3.4. Let (W, S) , Φ and $\mathbb{A}$ be as in 3.1. To each $\alpha \in \Phi$ we associate the reflection s_α given by

$$s_\alpha(v) = v - 2(v \cdot \alpha)\alpha$$

for each $v \in \mathbb{A}$. A *wall* of Φ is the fixed point set of one of these reflection. A *Weyl chamber* of Φ is a connected component of $\mathbb{A}$ with all the walls removed. We call

the closure of a Weyl chamber a *sector* of Φ . The sectors are polyhedral cones; a *face* of Φ is a face of one of these cones. The set of all faces of Φ forms a simplicial complex called the *Coxeter complex* of (W, S) (or Φ). We will denote this simplicial complex by $\Sigma(\Phi)$. To each element α of Φ we associate the half-space

$$H_\alpha := \{v \in \mathbb{A} \mid v \cdot \alpha \geq 0\}.$$

A *root* of the Coxeter complex is the set of faces contained in one of these half-spaces. The map $\alpha \mapsto H_\alpha$ thus gives rise to a canonical bijection from Φ to the set of roots of $\Sigma(\Phi)$.

Conventions 3.5. Let Δ be an irreducible spherical building. Then every apartment of Δ is isomorphic to the Coxeter complex $\Sigma(\Phi)$ for some Φ as in 3.1. The corresponding Coxeter system (W, S) is usually called the *type* of Δ . We prefer, instead, to say that Δ is *of type* Φ . (Thus a building of type B_ℓ is the same thing as building of type C_ℓ .) Let Σ be an apartment of Δ . A *root* of Σ is the image of a root of $\Sigma(\Phi)$ under an isomorphism from $\Sigma(\Phi)$ to Σ . Thus for each such isomorphism we have a canonical bijection from Φ to the set of roots of Σ . We will usually assume that an isomorphism from $\Sigma(\Phi)$ to Σ is fixed and identify Φ with the set of roots of Σ via this bijection.

Definition 3.6. Let Δ be an irreducible spherical building of rank at least two. For each root α of Δ (i.e. of some apartment of Δ), let U_α be the intersection of the stabilizers in $\text{Aut}(\Delta)$ of all the chambers that are contained in some panel of Δ which contains two chambers in α . It follows from Corollary 3.14 in [14] that U_α acts trivially on the set of chambers in α itself. (The subgroups of the form U_α are called the *root groups* of Δ .) The building Δ is *Moufang* (equivalently, “ Δ satisfies the *Moufang condition*”) if the following hold:

- (i) Δ is thick (i.e. every panel contains at least three chambers).
- (ii) For each root α of Δ , the root group U_α acts transitively on the set of apartments containing α .

Tits showed that thick irreducible spherical buildings of rank at least three, as well as all the irreducible residues of rank two of such a building, *always* satisfy the Moufang condition; see, for example, Theorems 11.6 and 11.8 in [14] for a proof. Irreducible spherical buildings of rank at least two satisfying the Moufang condition were classified in [10] and [13]. See [15, 30.14] for a summary of the results.

Proposition 3.7. *Let Δ be an irreducible spherical building of rank $\ell \geq 2$ satisfying the Moufang property. Then for each root α of Δ , the root group U_α acts sharply transitively on the set of apartments containing α .*

Proof. This is proved, for example, in Theorem 9.3 and Proposition 11.4 of [14]. $\square$

Now suppose that Δ is a building of rank one. In other words, Δ is simply a set (whose elements are the chambers of Δ) and $\text{Aut}(\Delta)$ is the full symmetric group on Δ . The apartments of Δ are the two-element subsets of Δ and thus the roots of Δ are just the one-element subsets of Δ . Normally we will use letters like x and y to name elements of Δ ; when we want to emphasize that an element of Δ is being considered as a root, however, we will give it a name like α or β .

Definition 3.8. Let Δ be a building of rank one, i.e. of type $\Phi := A_1$, let Σ be an apartment of Δ and let the two elements of Σ be identified with the two elements

of Φ . A *Moufang structure* on Δ is a collection

$$(U_\alpha)_{\alpha \in \Phi}$$

of non-trivial subgroups of $\text{Aut}(\Delta)$ such that the following hold:

- (i) For each $\alpha \in \Phi$, the subgroup U_α fixes α and acts sharply transitively on $\Delta \setminus \{\alpha\}$; and
- (ii) For each $\alpha \in \Phi$, the subgroup U_α is normalized by the stabilizer of α in the group $G := \langle U_\alpha, U_{-\alpha} \rangle$.

The groups U_α^g for all $\alpha \in \Phi$ and all $g \in G$ are called *root groups*. A Moufang structure on Δ is independent of the choice of Σ and its identification with Φ up to conjugation in the group G . Since the root groups are required to be non-trivial, Δ can have a Moufang structure only if it is thick, i.e. if $|\Delta| \geq 3$. When we say that Δ is Moufang, we mean that we have a particular Moufang structure on Δ in mind (whose root groups we will always call U_α, U_β , etc.). Note that with this convention, 3.7 holds also when $\ell = 1$ (with “satisfying the Moufang property” interpreted as meaning “having a Moufang structure with root groups U_α ”).

Remark 3.9. Let Δ be an irreducible spherical building of rank $\ell \geq 2$ which satisfies the Moufang condition and let P be a panel of Δ viewed as a set of chambers. For each chamber x in P , let α be an arbitrary root of Δ containing x but no other chamber in P . By 3.7, U_α acts faithfully on P . Furthermore, the permutation group induced by the root group U_α on P is independent of the choice of α . (This follows from Proposition 11.11 in [14].) Hence every rank one residue of Δ inherits a canonical Moufang structure from Δ whose root groups are isomorphic to root groups of Δ .

Proposition 3.10. *Let Δ be an irreducible spherical building of type Φ satisfying the Moufang condition as defined in 3.6 and 3.8 and let Σ be an apartment of Δ (to which we apply 3.5). Then for each $\alpha \in \Phi$ (i.e. to each root α of Σ), the following hold:*

- (i) *There exist maps λ and κ from U_α^* to $U_{-\alpha}^*$ such that for each $u \in U_\alpha^*$, the product*

$$m_\Sigma(u) := \kappa(u)u\lambda(u)$$

maps Σ to itself and induces the unique reflection s_α defined in 3.4 on Φ (which interchanges the roots α and $-\alpha$).

- (ii) $m_\Sigma(u)^{-1} = m_\Sigma(u^{-1})$ for each $u \in U_\alpha$.
- (iii) $m_\Sigma(\kappa(u)) = m_\Sigma(\lambda(u)) = m_\Sigma(u)$ for all $u \in U_\alpha$.

Proof. The first assertion is a consequence of 3.7 and the other two follow from the first; see, for example, in [13, 6.1–6.3]. $\square$

Proposition 3.11. *Let Δ be an irreducible spherical building satisfying the Moufang condition as defined in 3.6 and 3.8 and let $G^\dagger$ be the subgroup of $\text{Aut}(\Delta)$ generated by all the root groups of Δ . Then $G^\dagger$ acts transitively on the set of all pairs (Σ, C) , where Σ is an apartment of Δ and C is a chamber of Σ .*

Proof. This is proved, for example, in Proposition 11.12 of [14]. $\square$

Note in the rank one case, 3.11 just says that $G^\dagger$ is a 2-transitive permutation group on Δ .

Definition 3.12. Let Δ be an irreducible spherical building of type Φ satisfying the Moufang condition and let Σ be an apartment of Δ (to which we apply 3.5). For each $\alpha \in \Phi$, let U_α be the corresponding root group. The *root datum* of Δ (based at Σ) is the pair $(\Sigma, (U_\alpha)_{\alpha \in \Phi})$.

Let Δ be as in 3.12. By 3.11, the root datum of Δ is, up to conjugation in the group $G^\dagger$, independent of the choice of the apartment Σ . Note that a root datum and a Moufang structure on Δ (as defined in 3.8) are essentially the same thing when Δ has rank one. By [13, 40.17], Δ is uniquely determined by its root datum when the rank of Δ is at least two.

Theorem 3.13. *Let Δ be an irreducible spherical building of type Φ satisfying the Moufang condition, let the notion of the interval from one element of Φ to another be as in 3.2 and let Σ be an apartment of Δ (to which we apply 3.5). Then for all ordered pairs α, β of elements of Φ such that $\beta \neq \pm\alpha$,*

$$[U_\alpha, U_\beta] \subset U_{\gamma_1} U_{\gamma_2} \cdots U_{\gamma_s},$$

where $(\gamma_1, \gamma_2, \dots, \gamma_s)$ is the interval (α, β) , if the interval (α, β) is not empty and $[U_\alpha, U_\beta] = 1$ if it is.

Proof. This is proved in [7, 6.12(ii)]. $\square$

Definition 3.14. Let Δ be an irreducible spherical building of type Φ satisfying the Moufang condition, let Σ be an apartment of Δ (to which we apply 3.5) and let

$$(\Sigma, (U_\alpha)_{\alpha \in \Phi})$$

be the root datum of Δ based at Σ as defined in 3.12. A *valuation* of this root datum is a collection $\varphi := (\varphi_\alpha)_{\alpha \in \Phi}$ of non-constant maps φ_α from U_α^* to $\mathbb{R}$ such that the following hold:

V1: For each $\alpha \in \Phi$ and each $k \in \mathbb{R}$, the set

$$U_{\alpha,k} := \{u \in U_\alpha \mid \varphi_\alpha(u) \geq k\}$$

is a subgroup of U_α , where we assign $\varphi_\alpha(1)$ the value ∞ (so that $1 \in U_{\alpha,k}$ for all k).

V2: For all $\alpha, \beta \in \Phi$ such that $\alpha \neq \pm\beta$ and for all $k, l \in \mathbb{R}$,

$$[U_{\alpha,k}, U_{\beta,l}] = U_{\gamma_1, p_1 k + q_1 l} U_{\gamma_2, p_2 k + q_2 l} \cdots U_{\gamma_s, p_s k + q_s l},$$

where γ_i, p_i, q_i and s are as in 3.3.

V3: For all $\alpha, \beta \in \Phi$, all $u \in U_\alpha^*$ and all $g \in U_\beta$, the quantity

$$t := \varphi_{s_\alpha(\beta)}(g^{m_\Sigma(u)}) - \varphi_\beta(g)$$

is independent of g and if $\alpha = \beta$, then $t = -2\varphi_\alpha(u)$. Here s_α is as in 3.4 and $m_\Sigma(u)$ is as in 3.10.i.

Note that the condition (V2) is vacuous when the rank of Δ is one.

Definition 3.15. Let Δ, Φ and Σ be as in 3.14 and suppose that φ and ψ are two valuations of the root datum of Δ based at Σ . Then φ and ψ are *equipollent* if for some x in the ambient Euclidean space $\mathbb{A}$ of the root system Φ ,

$$\varphi_\alpha(u) = \psi_\alpha(u) + \alpha \cdot x$$

for all $\alpha \in \Phi$ and all $u \in U_\alpha^*$ (in which case we write $\varphi = \psi + x$).

Proposition 3.16. *Let Δ , Φ and Σ be as in 3.14 and suppose that φ and ψ are two valuations of the root datum of Δ based at Σ such that $\varphi_\alpha = \psi_\alpha$ for some $\alpha \in \Phi$. Then ψ and φ are equipollent (as defined in 3.15).*

Proof. This holds by Proposition 6 in [12]. For more details, see Theorem 3.41 in [15]. $\square$

Definition 3.17. Let Δ , Φ and Σ be as in 3.14, let $\varphi = (\varphi_\alpha)_{\alpha \in \Phi}$ be a valuation of the root datum of Δ based at Σ and let ρ be an automorphism of Δ mapping Σ to itself. We will say that φ is ρ -invariant if

$$\varphi_\alpha(u) = \varphi_{\alpha^\rho}(u^\rho)$$

for all $\alpha \in \Phi$ and all $u \in U_\alpha^*$.

Proposition 3.18. *Let Δ , Φ and Σ be as in 3.14, let φ be a valuation of the root datum of Δ based at Σ , let w be an element of the root group U_α^* such that $\varphi_\alpha(w) = 0$ and let $m_0 = m_\Sigma(w)$. Then for each $\alpha \in \Phi$,*

$$\varphi_\alpha(g^{m_0 m_\Sigma(u)}) - \varphi_\alpha(g) = 2\varphi_\alpha(u)$$

for all $g, u \in U_\alpha^*$.

Proof. Let β be the root opposite α in Σ . Let $g, u \in U_\alpha^*$ and let $v = \kappa(u) \in U_\beta^*$, where κ is as in 3.10.i. Then $m_\Sigma(u) = m_\Sigma(v)$ by Proposition 11.24 in [14], $\varphi_\beta(v) = -\varphi_\alpha(u)$ by Proposition 3.25 in [15] (or [7, 10.10]) and

$$\varphi_\alpha(y^{m_\Sigma(v)}) = \varphi_\beta(y) - 2\varphi_\beta(v)$$

for all $y \in U_\beta$ by condition (V3) (with both roots equal to β). Thus

$$\begin{aligned} \varphi_\alpha(g^{m_0 m_\Sigma(u)}) &= \varphi_\alpha(g^{m_0 m_\Sigma(v)}) \\ &= \varphi_\beta(g^{m_0}) - 2\varphi_\beta(v) \\ &= \varphi_\beta(g^{m_0}) + 2\varphi_\alpha(u) \end{aligned}$$

By another application of condition (V3) (this time with both roots equal to α) and the choice of w , we have

$$\varphi_\beta(g^{m_0}) = \varphi_\alpha(g).$$

$\square$

4. NON-DISCRETE EUCLIDEAN BUILDINGS

In this section we assemble a few basic facts about non-discrete Euclidean buildings. We start with the definition.

Notation 4.1. Let W , $\mathbb{A}$ and Φ be as in 3.1. Let $\mathbb{W}$ denote the group generated by W and the group T consisting of all translations of $\mathbb{A}$. Thus $\mathbb{W}$ is a group of isometries of $\mathbb{A}$. Moreover, $\mathbb{W} = TW$ and T is a normal subgroup of $\mathbb{W}$.

Definition 4.2. Let Φ and $(\mathbb{A}, \mathbb{W})$ be as in 4.1 and let H_α for all $\alpha \in \Phi$ be as in 3.4. Let X be a set and let $\mathcal{A}$ be a family of injections of $\mathbb{A}$ into X . The elements of $\mathcal{A}$ will be called *charts* and the images of charts will be called *apartments*. Sets of the form $f(H_\alpha)$ for some chart f and some $\alpha \in \Phi$ will be called *roots* of $(X, \mathcal{A})$ and sets of the form $f(S)$ for some chart f and some sector (respectively, face) of Φ (as defined in 3.4) will be called *sectors* (respectively, *faces*) of $(X, \mathcal{A})$. The pair

$(X, \mathcal{A})$ is a *non-discrete Euclidean building of type Φ* if the following six axioms hold:

- A1:** If $f \in \mathcal{A}$ and $w \in \mathbb{W}$, then $f \circ w \in \mathcal{A}$.
- A2:** If $f, f' \in \mathcal{A}$, then the set

$$M := \{v \in \mathbb{A} \mid f(v) \in f'(\mathbb{A})\}$$

is closed and convex and there exists $w \in \mathbb{W}$ such that the maps f and $f' \circ w$ coincide on M .

- A3:** Every two points of X are contained in a common apartment.
- A4:** If S and S' are sectors of $(X, \mathcal{A})$, then there exists an apartment containing sectors S_1 and S'_1 such that $S_1 \subset S$ and $S'_1 \subset S'$.
- A5:** Three apartments which intersect pairwise in roots have a non-empty intersection.
- A6:** There is a metric d on X such that for all $v, z \in \mathbb{A}$ and all $f \in \mathcal{A}$, $d(f(v), f(z))$ equals the Euclidean distance between v and z .

If $(X, \mathcal{A})$ is a non-discrete Euclidean building of type Φ , we call the pair $(\mathbb{A}, \mathbb{W})$, which is uniquely determined by Φ , its *model* and we define the *dimension* of $(X, \mathcal{A})$ to be the dimension of $\mathbb{A}$. By (A3), the metric d in (A6) is unique.

Various equivalent definitions of a non-discrete Euclidean building (and a proof of their equivalence) can be found in Theorem 1.21 of [6]. See also Proposition 2.21 of [6].

Definition 4.3. Let $(X, \mathcal{A})$ and $(X', \mathcal{A}')$ be two non-discrete Euclidean buildings having the same type Φ . An *isomorphism* ψ from $(X, \mathcal{A})$ to $(X', \mathcal{A}')$ is a bijection from X to X' such that

$$\mathcal{A}' = \{\psi \circ f \mid f \in \mathcal{A}\}.$$

We denote by $\text{Aut}(X, \mathcal{A})$ the group of all isomorphisms from $(X, \mathcal{A})$ to itself.

Definition 4.4. Let $(X, \mathcal{A})$ be a non-discrete Euclidean building of type Φ , let $(\mathbb{A}, \mathbb{W})$ be the model of $(X, \mathcal{A})$, let τ be an isometry of $\mathbb{A}$ normalizing the group $\mathbb{W}$ and let

$$\mathcal{A}_\tau = \{f \circ \tau \mid f \in \mathcal{A}\}.$$

Then $(X, \mathcal{A}_\tau)$ is a non-discrete Euclidean building of type Φ with the same underlying metric structure as $(X, \mathcal{A})$. Now suppose in addition that $\tau^2 = 1$. Then an isomorphism from $(X, \mathcal{A})$ to $(X, \mathcal{A}_\tau)$ is automatically an isomorphism from $(X, \mathcal{A}_\tau)$ to $(X, \mathcal{A})$. We call such an isomorphism a τ -*automorphism* of $(X, \mathcal{A})$.

Definition 4.5. We say that two non-discrete Euclidean buildings $(X, \mathcal{A})$ and $(X', \mathcal{A}')$ are *equivalent* (or one is a *dilation* of the other) if they have the same type Φ and therefore the same model $(\mathbb{A}, \mathbb{W})$, $X = X'$ and

$$\mathcal{A}' = \{f \circ \delta \mid f \in \mathcal{A}\}$$

for some dilation δ of $\mathbb{A}$ (where *dilation* means multiplication by a non-zero constant). Thus equivalent non-discrete Euclidean buildings have the same underlying metric structure up to a constant positive factor.

Definition 4.6. Let $(X, \mathcal{A})$ be a non-discrete Euclidean building with model $(\mathbb{A}, \mathbb{W})$ and let $x, x' \in X$. By (A3), there exists a chart f and points $x_1, x'_1 \in \mathbb{A}$ such that $f(x_1) = x$ and $f(x'_1) = x'$. The *interval* $[x, x']$ is the image under f of the interval $[x_1, x'_1]$. By (A2), the interval $[x, x']$ is independent of the chart f . By the CAT(0)

property (proved, for example, in Proposition 2.10 of [5]), the interval $[x, x']$ is, in fact, the unique geodesic connecting x to x' .

Notation 4.7. Let $(X, \mathcal{A})$ be a non-discrete Euclidean building of type Φ . Two faces F and F' of $(X, \mathcal{A})$ (as defined in 4.2) are called *parallel* if they are at finite Hausdorff distance, i.e. if both

$$\sup_{x' \in F'} d(x', F)$$

and

$$\sup_{x \in F} d(x, F')$$

are finite. By (A6), this is an equivalence relation on faces. For each face F of $(X, \mathcal{A})$, we denote by F^∞ the corresponding parallel class and for each apartment A of $(X, \mathcal{A})$, we denote by A^∞ the set of parallel classes of faces containing a face of A . If b and b' are two parallel classes, we set $b \leq b'$ whenever

$$\sup_{x' \in F'} d(x', F) < \infty.$$

for all $F \in b$ and all $F' \in b'$. This notion makes the set of parallel classes of faces into a simplicial complex. We denote this simplicial complex by $(X, \mathcal{A})^\infty$. By Proposition 1 in [12] (see also Property 1.7 in [6]), $(X, \mathcal{A})^\infty$ is, in fact, a spherical building of type Φ (as defined in 3.4) and the map $A \mapsto A^\infty$ is a bijection from the set of apartments of $(X, \mathcal{A})$ to the set of apartments of $(X, \mathcal{A})^\infty$. The building $(X, \mathcal{A})^\infty$ is called the *building at infinity* of $(X, \mathcal{A})$. It is irreducible (since it is of type Φ) and its rank is the same as the dimension of $(X, \mathcal{A})$.

Notation 4.8. Let $(X, \mathcal{A})$ be a non-discrete Euclidean building of type Φ with model $(\mathbb{A}, \mathbb{W})$, let A be an apartment of $(X, \mathcal{A})$, let x be a point of A and let $\Sigma = A^\infty$. Let $\Sigma(\Phi)$ and H_α (for each $\alpha \in \Phi$) be as in 3.4 and suppose that Φ is identified with the set of roots of Σ via an isomorphism from $\Sigma(\Phi)$ to Σ as described in 3.5. Then there exists a unique chart f such that the following hold:

- (i) $f(\mathbb{A}) = A$.
- (ii) $f(0) = x$.
- (iii) For each $\alpha \in \Phi$, a sector S of Φ is contained in the half-space H_α if and only if the chamber $f(S)^\infty$ of Σ is contained in the root α of Σ .

Let $f_{A,x}$ denote the chart f .

Remark 4.9. Let Φ and $(\mathbb{A}, \mathbb{W})$ be as in 4.1. Then $(\mathbb{A}, \mathbb{W})$ is a non-discrete Euclidean building of type Φ with just one apartment. It thus has a building at infinity whose faces are of the form F^∞ for some face F of Φ .

Notation 4.10. Let $(X, \mathcal{A})$ be a non-discrete Euclidean building of type Φ and let $(\mathbb{A}, \mathbb{W})$ be its model. Let F be a face of $(X, \mathcal{A})$. Then F is the image of a face of Φ (as defined in 3.4) under some chart f . The *vertex* of F is the image of the origin of $\mathbb{A}$ under f ; by (A2), this notion is independent of the choice of f . Now let x be a point of X . We declare two points y and z of $X \setminus \{x\}$ to be *equivalent at x* if

$$[x, y] \cap B = [x, z] \cap B$$

for some open ball B centered at x , where $[x, y]$ and $[x, z]$ are intervals as defined in 4.6. This is an equivalence relation on $X \setminus \{x\}$. For each $y \in X \setminus \{x\}$, let $g_x(y)$ denote its equivalence class. We declare two faces F and F_1 with vertex x to be

equivalent if $g_x(F) = g_x(F_1)$. By (A2), this holds if and only if $F \cap B = F_1 \cap B$ for some open ball B centered at x . The equivalence class of a face F is called the *germ of F* and a *germ at x* is the germ of a face with vertex x . The set of germs at x form a simplicial complex on the set $g_x(X \setminus \{x\})$. As observed in Section 1.3 of [6], this simplicial complex is a building of type Φ whose apartments are the sets $g_x(A)$ for all apartments A of $(X, \mathcal{A})$ containing x . We call this building the *residue of $(\Delta, \mathcal{A})$ at x* and denote it by $(\Delta, \mathcal{A})_x$. (The residue at x is called the *building of directions* at x in [6].)

The residues of a non-discrete Euclidean building $(X, \mathcal{A})$ are not necessarily thick, and the residues at different points of X are not necessarily isomorphic to each other. They are all, however, of type Φ .

Notation 4.11. Let $(X, \mathcal{A})$ be a non-discrete Euclidean building, let $x \in X$, let the residue $(X, \mathcal{A})_x$ of $(X, \mathcal{A})$ at x be as in 4.10 and let f be an element of $\mathcal{A}$ mapping the origin 0 to x . Then there exists a unique type-preserving isomorphism from the Coxeter complex $\Sigma(\Phi)$ (as defined in 3.4) to an apartment of the residue $(X, \mathcal{A})_x$ which maps each face F of Φ to the germ at x containing $f(F)$. We denote this isomorphism by f_* .

For the rest of this section, we examine the special case that the building at infinity of a non-discrete Euclidean building is Moufang.

Theorem 4.12. *Let $(X, \mathcal{A})$ be a non-discrete Euclidean building of type Φ with model $(\mathbb{A}, \mathbb{W})$ such that the building at infinity $\Delta := (X, \mathcal{A})^\infty$ satisfies the Moufang property as defined in 3.6, let ℓ denote the dimension of $(X, \mathcal{A})$, let A be an apartment of $(X, \mathcal{A})$, let $\Sigma = A^\infty$, let x be a point of A , let $\Sigma(\Phi)$ be as in 3.4 and let*

$$H_{\alpha, k} = \{v \in \mathbb{A} \mid v \cdot \alpha \geq k\}$$

for all $\alpha \in \Phi$ and all $k \in \mathbb{R}$. Let Φ be identified with the set of roots of Σ via an isomorphism from $\Sigma(\Phi)$ to Σ as described in 3.5 and let $f := f_{A, x}$ be as in 4.8. If $\ell \geq 2$, then the following hold:

- (i) *For every root $\alpha \in \Phi$, there exists a canonical injection from the root group U_α into $\text{Aut}(X, \mathcal{A})$ such that for each $u \in U_\alpha$, its image under this injection induces u on Δ .*
- (ii) *For every $\alpha \in \Phi$, there exists a map φ_α from the root group U_α^* of Δ to $\mathbb{R}$ such that*

$$\text{Fix}_A(u) = A \cap A^u = f(H_{\alpha, \varphi_\alpha(u)})$$

for each $u \in U_\alpha^$.*

Proof. See Sections 10 and 11 of [12]. □

Note that under the hypotheses of 4.12, we always identify each root group U_α with its image under the injection in 4.12.i. Thus, in particular, the u in 4.12.ii is really the canonical image in $\text{Aut}(X, \mathcal{A})$ of an element $u \in U_\alpha$.

With the following definition we describe those non-discrete Euclidean buildings which were studied in [2].

Definition 4.13. A *Bruhat-Tits space* is a non-discrete Euclidean building $(X, \mathcal{A})$ such that the following hold:

- (i) The spherical building $\Delta := (X, \mathcal{A})^\infty$ is Moufang (in the sense of 3.6 or 3.8).

(ii) The conclusions of 4.12 hold.

Let $(X, \mathcal{A})$ be a non-discrete Euclidean building satisfying (i) and let ℓ denote the dimension of $(X, \mathcal{A})$. If $\ell = 1$, saying that Δ is Moufang means we have a particular Moufang structure on Δ in mind, and (ii) is to be interpreted with respect to this Moufang structure. If $\ell \geq 2$, then (ii) holds automatically.

A non-discrete Euclidean building of rank $\ell \geq 3$ always satisfies 4.13.i. (This is proved, for example, [13, 40.3].) Thus in dimension three or higher, “non-discrete Euclidean building” and “Bruhat-Tits space” are the same thing.

Convention 4.14. Let Δ be a Moufang building of rank one in the sense of 3.8. When we say that “ Δ is the building at infinity of the Bruhat-Tits space $(X, \mathcal{A})$,” we mean that the conclusions of 4.12 hold with respect to the particular Moufang structure on Δ we have in mind.

The following results of Bruhat-Tits and Tits are fundamental.

Theorem 4.15. *Let $(X, \mathcal{A})$, A , x , Δ , Σ and φ_α for $\alpha \in \Phi$ be as in 4.12. Then*

$$\varphi := \{\varphi_\alpha \mid \alpha \in \Phi\}$$

is a valuation of the root datum of Δ based at Σ . Moreover, the valuation φ is independent of the choice of the point x in A up to equipollence (as defined in 3.15).

Proof. This is the first part of Theorem 3 in [12]. □

Theorem 4.16. *Let Δ be an irreducible spherical building of type Φ satisfying the Moufang condition, let Σ be an apartment of Δ (to which 3.5 is applied) and let φ be a valuation of the root datum of Δ based at Σ as defined in 3.14. Then there exists a Bruhat-Tits space $(X, \mathcal{A})$ of type Φ , an apartment A of $(X, \mathcal{A})$ and a point x_A of A such that the following hold:*

- (i) Δ is the building at infinity of $(X, \mathcal{A})^\infty$ (in the sense of 4.14 if the rank of Δ is one) and $\Sigma = A^\infty$.
- (ii) For each $\alpha \in \Phi$, φ_α is the map which appears in 4.12.ii when 4.12 is applied to the triple $(X, \mathcal{A})$, A and x_A .

If $(X', \mathcal{A}')$, A' and x'_A is a second triple with these properties, then there exists an isomorphism from $(X, \mathcal{A})$ to $(X', \mathcal{A}')$ mapping A to A' and x_A to x'_A .

Proof. Existence is proved in Section 7.4 of [2] and uniqueness in Proposition 6 of [12]. □

5. THE REE AND SUZUKI GROUPS

In this section we collect a few well known facts about the Ree and Suzuki groups. All of these results are contained more or less explicitly in [9] and [11].

There are three families of Ree and Suzuki groups. Beginning in 5.1 and for the rest of this paper, we will refer to three cases which we will call “case B,” “case F” and “case G.”

Notation 5.1. Let K be a field of characteristic positive p and suppose that θ is a *Tits endomorphism* of K . This means that θ is an endomorphism of K such that θ^2 is the Frobenius map $x \mapsto x^p$. Thus, in particular, $F := K^\theta$ is a subfield of K isomorphic to K which contains the subfield K^p . Suppose, too, that $p = 2$ in

cases B and F and $p = 3$ in case G. In case B let L be an additive subgroup of K containing F such that $L \cdot F \subset L$ (so L is a vector space over F) and $K = \langle L \rangle$ (where $\langle L \rangle$ denotes the subring of K generated by L) and let Λ denote the indifferent set (K, L, L^θ) as defined in [13, 10.1]. In case F let Λ denote the composition algebra (K, F) as defined in [15, 30.17]. In case G let Λ denote the hexagonal system $(K/F)^\circ$ as defined in [13, 15.20]. Let Δ denote the building $B_2^{\mathcal{D}}(\Lambda)$ in case B, the building $F_4(\Lambda)$ in case F and the building $G_2(\Lambda)$ in case G in the notation described in [15, 30.17]. Thus $B_2^{\mathcal{D}}(\Lambda)$ is the Moufang quadrangle called $\mathcal{Q}_{\mathcal{D}}(\Lambda)$ in [13, 16.4] and $G_2(\Lambda)$ the Moufang hexagon called $\mathcal{H}(\Lambda)$ in [13, 16.8]. The type Φ of the building Δ is B_2 in case B, F_4 in case F and G_2 in case G. (Alternatively, we can define Δ to be the unique building of type Φ whose root datum is as described in 5.3 below.)

The building Δ in 5.1 is split if and only if K is perfect; if K is not perfect, Δ is simply a building of mixed type (as defined, for example, in [15, 30.24]).

The following element τ plays a central role from now on.

Notation 5.2. Let Φ be as in 5.1, let $\mathbb{A}$ be the ambient space of Φ and let S be a sector of Φ . There is a unique non-trivial element of $\text{Aut}(\Phi)$ (as defined in 3.1) fixing S . This automorphism induces a non-type-preserving automorphism of the Coxeter complex $\Sigma(\Phi)$ and has order two. We denote it by τ .

Theorem 5.3. *Let Δ , Φ , K , L , etc. be as in 5.1, let τ and S be as in 5.2, let Σ be an apartment of Δ , let C be a chamber of Σ , let ψ be the unique special isomorphism from $\Sigma(\Phi)$ to Σ mapping S to C and let Φ be identified with the set of roots of Σ via ψ as indicated in 3.5. Then for each $\alpha \in \Phi$, there exists an isomorphism x_α from the additive group of L in case B, respectively, the additive group of K in cases F and G, to the root group U_α of Δ such that the collection $(x_\alpha)_{\alpha \in \Phi}$ has the following properties:*

- (i) *There exists a unique automorphism ρ of Δ mapping the pair (C, Σ) to itself such that*

$$x_\alpha(t)^\rho = x_{\tau(\alpha)}(t)$$

for all $\alpha \in \Phi$ and all $t \in K$ (or all $t \in L$).

- (ii) *In cases B and F,*

$$[U_\alpha, U_\beta] = 1$$

whenever $\angle(\alpha, \beta) \leq 90^\circ$,

$$[x_\alpha(s), x_\beta(t)] = x_{\alpha+\beta}(st)$$

for all $s, t \in K$ whenever $\angle(\alpha, \beta) = 120^\circ$ and

$$[x_\alpha(s), x_\beta(t)] = x_{\sqrt{2}\alpha+\beta}(s^\theta t)x_{\alpha+\sqrt{2}\beta}(st^\theta)$$

for all $s, t \in K$ (or all $s, t \in L$) whenever $\angle(\alpha, \beta) = 135^\circ$.

- (iii) *In case G, there exists parameters $\epsilon, \epsilon_1, \dots, \epsilon_4$ such that*

$$[U_\alpha, U_\beta] = 1$$

whenever $\angle(\alpha, \beta) \leq 90^\circ$,

$$[x_\alpha(s), x_\beta(t)] = x_{\alpha+\beta}(\epsilon st)$$

for all $s, t \in K$ whenever $\angle(\alpha, \beta) = 120^\circ$ and

$$[x_\alpha(s), x_\beta(t)] = x_{\sqrt{3}\alpha+\beta}(\epsilon_1 s^\theta t) x_{2\alpha+\sqrt{3}\beta}(\epsilon_2 s^2 t^\theta) \cdot x_{\sqrt{3}\alpha+2\beta}(\epsilon_3 s^\theta t^2) x_{\alpha+\sqrt{3}\beta}(\epsilon_4 s t^\theta)$$

for all $s, t \in K$ whenever $\angle(\alpha, \beta) = 150^\circ$. The parameters $\epsilon, \epsilon_1, \dots, \epsilon_4$ are all equal to $+1$ or -1 ; their values depend on the ordered pair (α, β) but not on s or t ; and their values are as in 5.4–5.6 if α, β both contain the chamber C .

(iv) In all three cases,

$$x_\beta(t)^{m_\Sigma(x_\alpha(1))} = x_{s_\alpha(\beta)}(\pm t)$$

for all $\alpha, \beta \in \Phi$ and all $t \in K$ (or L).

Proof. Suppose first that we are in case G and let the roots of Φ be numbered $\alpha_1, \dots, \alpha_{12}$ going around the origin clockwise, where the indices are to be read modulo 12. The set of roots of Σ can be identified with Φ so that α_i contains the chamber C if and only if $i \in [1, 6]$. Thus $\tau(\alpha_i) = \alpha_{7-i}$ for all i . By [13, 16.8], $[U_{\alpha_i}, U_{\alpha_j}] = 1$ whenever $i, j \in [1, 6]$ and $i < j \leq i+3$ and there exist isomorphisms x_i for each $i \in [1, 6]$ from the additive group of K to the root group U_{α_i} such that for all $s, t \in K$,

$$(5.4) \quad [x_1(s), x_6(t)] = x_2(-s^\theta t) x_3(-s^2 t^\theta) x_4(s^\theta t^2) x_5(st^\theta),$$

as well as

$$(5.5) \quad [x_1(s), x_5(t)] = x_3(-st)$$

and

$$(5.6) \quad [x_2(s), x_6(t)] = x_4(st).$$

By [13, 7.5], there exists a unique automorphism ρ of Δ mapping (C, Σ) to itself such that

$$x_i(t)^\rho = x_{7-i}(t)$$

for all $i \in [1, 6]$ and for all $t \in K$. Let H be the chamberwise stabilizer of Σ in $\text{Aut}(\Delta)$ and let Q be the subgroup of H consisting of those elements g such that for each root α of Σ , either g centralizes U_α or g inverts every element of U_α . Let

$$m_i = m_\Sigma(x_i(1))$$

for $i = 1$ and 6 and let

$$N = \langle m_1, m_6 \rangle.$$

By [11, 2.9(6)] and [13, 29.12], $|Q| = 4$ and the kernel of the action of N on Σ is $N \cap Q$ and $N/N \cap Q \cong D_{12}$. For each $i \in [7, 12]$, there thus is a unique shortest word g in m_1 and m_6 mapping α_j to α_i , where $j = 1$ if i is odd and $j = 6$ if i is even. We set $x_i(t) = x_j(t)^g$ for all $t \in K$. By [13, 6.2], $m_1^\rho = m_6$ and $m_6^\rho = m_1$. It follows that $x_i(t)^\rho = x_{\tau(i)}(t)$ for all $t \in K$ and for all $i \in [7, 12]$. Thus (i) holds. Let $\alpha, \beta \in \Phi$. By [13, 6.4] (or [11, 2.9]), there exists $g \in N$ mapping α to β such that $x_\alpha(t)^g = x_\beta(\pm t)$ for all $t \in K^*$. Since g is unique up to an element of $N \cap Q$, it follows that $x_\alpha(t)^g = x_\beta(\pm t)$ for all $t \in K^*$ and for all $g \in N$ mapping α to β . Thus (iv) holds (in case G). By 5.4–5.6, it follows that also (iii) holds.

The proof in case B is virtually the same, except that we cite [13, 16.4] in place of [13, 16.8] and use the observation that this time $m_\Sigma(u)^2 = 1$ for all $\alpha \in \Phi$ and all $u \in U_\alpha^*$ by 3.10.ii (since all the root groups are of exponent two). In case F, the

proof in the previous two cases can be suitably modified (using [15, 32.14] in place of [13, 16.4]). We do not give the details, however, since the claims in case F also follow from the observations in [11, 1.1–1.2]. $\square$

Definition 5.7. Let Δ and ρ be as in 5.3, let Δ^ρ denote the set of chambers fixed by ρ and let $G^\dagger$ be as in 3.11. Let G be the group induced on Δ^ρ by the centralizer of ρ in $G^\dagger$. The group G is commonly called $\text{Suz}(K, \theta, L)$ in case B (these are the *Suzuki groups*), $\text{Ree}(K, \theta)$ in case G (these are the *Ree groups*) and ${}^2F_4(K, \theta)$ in case F. The groups ${}^2F_4(K, \theta)$ were also discovered by Ree and are also sometimes called Ree groups, but these groups are now more commonly associated with Tits due to his thorough study of them in [11]. The name $\text{Suz}(K, \theta, L)$ is usually abbreviated to $\text{Suz}(K, \theta)$ when $L = K$.

We use the rest of this section to prove a result (5.15) about valuations of the root datum of the building Δ .

Proposition 5.8. *Let Δ , Φ , Σ and $(x_\alpha)_{\alpha \in \Phi}$ be as in 5.3, let $\alpha, \beta \in \Phi$, let $t \in K$, let $u \in K^*$ (or $t \in L$ and $u \in L^*$ in case B) and let*

$$h(u) = m_\Sigma(x_\alpha(1))m_\Sigma(x_\alpha(u)).$$

Then the following hold:

(i) *In cases B and F,*

$$x_\beta(t)^{h(u)} = x_\beta(u^{-\theta}t)$$

if $\angle(\alpha, \beta) = 45^\circ$ and

$$x_\beta(t)^{h(u)} = x_\beta(u^\theta t)$$

if $\angle(\alpha, \beta) = 135^\circ$.

(ii) *In case G,*

$$x_\beta(t)^{h(u)} = x_\beta(u^\theta t)$$

if $\angle(\alpha, \beta) = 30^\circ$ and

$$x_\beta(t)^{h(u)} = x_\beta(u^{-\theta}t)$$

if $\angle(\alpha, \beta) = 150^\circ$.

(iii) *In all three cases,*

$$x_\beta(t)^{h(u)} = x_\beta(u^{-2}t)$$

if $\alpha = \beta$,

$$x_\beta(t)^{h(u)} = x_\beta(u^{-1}t)$$

if $\angle(\alpha, \beta) = 60^\circ$,

$$x_\beta(t)^{h(u)} = x_\beta(t)$$

if $\angle(\alpha, \beta) = 90^\circ$,

$$x_\beta(t)^{h(u)} = x_\beta(ut)$$

if $\angle(\alpha, \beta) = 120^\circ$ and

$$x_\beta(t)^{h(u)} = x_\beta(u^2t)$$

if $\alpha = -\beta$.

Proof. By 3.10.i, $h(u)$ acts trivially on Σ and hence normalizes U_β . The claims hold for $\beta \neq \pm\alpha$ by parts (ii) and (iii) of 5.3 and [11, 2.9].

Suppose that we can choose $\beta \in \Phi$ such that $\angle(\alpha, \beta) = 120^\circ$. By 5.3, we have

$$(5.9) \quad [x_\alpha(t), x_\beta(s)] = x_{\alpha+\beta}(\epsilon st)$$

and

$$(5.10) \quad [x_{\alpha+\beta}(s), x_{-\alpha}(t)] = x_\beta(\epsilon' st)$$

for all $s \in K$, where $\epsilon, \epsilon' = \pm 1$. We know that $x_\beta(1)^{h(u)} = x_\beta(u)$ and

$$x_{\alpha+\beta}(t)^{h(u)} = x_{\alpha+\beta}(u^{-1}t).$$

Setting $s = 1$ in 5.9 and conjugating by $h(u)$, we obtain

$$[x_\alpha(t)^{h(u)}, x_\beta(u)] = x_{\alpha+\beta}(\epsilon u^{-1}t).$$

Comparing this identity with 5.9 itself, we conclude that $x_\alpha(t)^{h(u)} = x_\alpha(u^{-2}t)$. Setting $s = 1$ in 5.10 and conjugating by $h(u)$, we conclude similarly that $x_{-\alpha}(t)^{h(u)} = x_{-\alpha}(u^2t)$.

Suppose next that there is no $\beta \in \Phi$ such that $\angle(\alpha, \beta) = 120^\circ$. Then we are in case B, so $\text{char}(K) = 2$, and we can find β such that $\angle(\alpha, \beta) = 135^\circ$. Thus

$$(5.11) \quad [x_\alpha(t), x_\beta(s)] = x_{\sqrt{2}\alpha+\beta}(t^\theta s) x_{\alpha+\sqrt{2}\beta}(ts^\theta)$$

and

$$(5.12) \quad [x_{\sqrt{2}\alpha+\beta}(s), x_{-\alpha}(t)] = x_{\alpha+\sqrt{2}\beta}(s^\theta t) x_\beta(st^\theta)$$

by 5.3.ii. We know that $h(u)$ centralizes U_3 , $x_\beta(1)^{h(u)} = x_\beta(t^\theta)$ and

$$x_{\sqrt{2}\alpha+\beta}(t^\theta)^{h(u)} = x_{\sqrt{2}\alpha+\beta}(u^{-\theta}t^\theta).$$

Setting $s = 1$ in 5.11 and conjugating by $h(u)$, we obtain

$$[x_\alpha(t)^{h(u)}, x_\beta(u^\theta)] = x_{\sqrt{2}\alpha+\beta}(u^{-\theta}t^\theta) x_{\alpha+\sqrt{2}\beta}(t).$$

Comparing this identity with 5.11 itself, we conclude again that

$$x_\alpha(t)^{h(u)} = x_\alpha(u^{-2}t).$$

Setting $s = 1$ in the identity 5.12 and conjugating by $h(u)$, we conclude similarly that $x_{-\alpha}(t)^{h(u)} = x_{-\alpha}(u^2t)$ also in this case. $\square$

Corollary 5.13. *Let Δ , Φ , Σ and $(x_\alpha)_{\alpha \in \Phi}$ be as in 5.3, let $G^\dagger$ be as in 3.11 and let H be the chamberwise stabilizer of Σ in $G^\dagger$. Then for each $\alpha \in \Phi$ and for each $h \in H$, there exists $z \in K^*$ (or $z \in F^*$ in case B, where $F = K^\theta$) such that*

$$x_\alpha(t)^h = x_\alpha(zt)$$

for all $t \in K$ (or all $t \in L$).

Proof. By [13, 33.9],

$$H = \langle m_\Sigma(x_\alpha(1))m_\Sigma(x_\alpha(u)) \mid \alpha \in \Phi \text{ and } u \in K^* \text{ (or } L^*) \rangle.$$

The claim holds, therefore, by 5.8. $\square$

Notation 5.14. Let Φ be as in 5.1. Let $\Phi = \Phi_0 \cup \Phi_1$ be the partition of Φ into two subsets Φ_0 and Φ_1 such that two elements of Φ are in the same subset if and only if they have the same length.

Proposition 5.15. *Let Δ , Φ , Σ , ρ and $(x_\alpha)_{\alpha \in \Phi}$ be as in 5.3, let Φ_0 and Φ_1 be as in 5.14, let ν be a (real valued) valuation of the field K and let $p = \text{char}(K)$. Let*

$$\varphi_\alpha(x_\alpha(t)) = \nu(t)$$

for all $\alpha \in \Phi_0$ and all $t \in K^$ (or all $t \in L^*$), let*

$$\varphi_\alpha(x_\alpha(t)) = \nu(t^\theta)/\sqrt{p}$$

for all $\alpha \in \Phi_1$ and all $t \in K^$ (or all $t \in L^*$) and let*

$$\varphi = (\varphi_\alpha)_{\alpha \in \Phi}.$$

Then the following hold:

- (i) φ is a valuation of the root datum of Δ based at Σ .
- (ii) φ is ρ -invariant (as defined in 3.17) if and only if ν is θ -invariant.

Proof. The collection φ satisfies condition (V1) in 3.14 by the definition of a valuation. By parts (ii) and (iii) of 5.3 and some calculation, φ satisfies condition (V2). Choose $\alpha, \beta \in \Phi$, $u \in U_\alpha^*$ and $g \in U_\beta^*$. Suppose first that $\alpha = \beta$. By 5.3.iv,

$$\varphi_{-\alpha}(g^{m_\Sigma(u)}) = \varphi_\alpha(g^{m_\Sigma(u)m_\Sigma(x_\alpha(1))^{-1}}).$$

By 3.10.ii, therefore

$$\varphi_{-\alpha}(g^{m_\Sigma(u)}) = \varphi_\alpha(g^{(m_\Sigma(x_\alpha(1))m_\Sigma(-u))^{-1}}).$$

Hence

$$\varphi_{-\alpha}(g^{m_\Sigma(u)}) - \varphi_\alpha(g) = -2\varphi_\alpha(u)$$

by 5.8.iii. By 5.3.iv and 5.8 (and a similar calculation), the quantity

$$\varphi_{s_\alpha(\beta)}(g^{m_\Sigma(u)}) - \varphi_\beta(g)$$

is independent of g whenever $\beta \neq \alpha$. Thus φ satisfies condition (V3). Hence (i) holds. By 5.3.i, φ is ρ -invariant if and only if $\nu(t^\theta)/\sqrt{p} = \nu(t)$ for all $t \in K^*$. Thus (ii) holds. $\square$

6. SPHERICAL BUILDINGS FOR THE REE AND SUZUKI GROUPS

In this section we show that the G -set Δ^ρ defined in 5.7 has the structure of a spherical building satisfying the Moufang condition whose root groups generate the group G .

Notation 6.1. Let $\dot{\Phi}$ and $\dot{\mathbb{A}}$ be the sets called Φ and $\mathbb{A}$ in 5.1 and 5.2. We then set $\mathbb{A}$ equal to the set of fixed points of the automorphism τ of $\dot{\Phi}$ introduced in 5.2 (which we continue to call τ).

Proposition 6.2. *Let $\dot{\mathbb{A}}$, $\dot{\Phi}$, τ and $\mathbb{A}$ be as in 6.1 and let $\ddot{\alpha} = \dot{\alpha} + \dot{\alpha}^\tau$ (so $\ddot{\alpha} \in \mathbb{A}$ since $\tau^2 = 1$) for each $\dot{\alpha} \in \dot{\mathbb{A}}$. Then the following hold:*

- (i) *If $\dot{\alpha} \in \dot{\Phi}$, then $\ddot{\alpha} \neq 0$.*
- (ii) *Let*

$$\Phi = \{\ddot{\alpha}/|\ddot{\alpha}| \mid \dot{\alpha} \in \dot{\Phi}\}.$$

Then Φ is, up to an isometry of $\mathbb{A}$, the root system A_1 in cases B and G and $I_2(8)$ (as defined in 3.1) in case F.

(iii) For each $\dot{\alpha} \in \dot{\Phi}$, the half-space

$$H_{\alpha} := \{v \in \mathbb{A} \mid v \cdot \ddot{\alpha} \geq 0\}$$

of $\Sigma(\Phi)$ (as defined in 3.4) is the intersection of the half-space

$$H_{\dot{\alpha}} := \{v \in \dot{\mathbb{A}} \mid v \cdot \dot{\alpha} \geq 0\}$$

of $\Sigma(\dot{\Phi})$ with $\mathbb{A}$.

(iv) The map $\dot{F} \mapsto \dot{F} \cap \mathbb{A}$ is an inclusion-preserving bijection from the set of τ -invariant faces of $\dot{\Phi}$ to the set of faces of Φ .

Proof. In cases B and G, the dimension of $\dot{\mathbb{A}}$ is only two; we leave it to the reader to verify all the claims in these two cases. In case F, these assertions are proved in Section 1.3 of [11]. $\square$

Lemma 6.3. *Let $\dot{\Phi}$ and τ be as in 6.1, let Φ be as in 6.2.ii, let $\dot{\Delta}$ be an arbitrary building of type $\dot{\Phi}$ which is not necessarily thick, let ρ be a non-type-preserving automorphism of $\dot{\Delta}$ of order two and let $\dot{\Sigma}$ be an apartment of $\dot{\Delta}$. Then the following are equivalent:*

- (i) $\dot{\psi} \circ \tau = \rho \circ \dot{\psi}$ for some type-preserving isomorphism $\dot{\psi}$ from $\Sigma(\dot{\Phi})$ to $\dot{\Sigma}$.
- (ii) ρ fixes two opposite chambers of $\dot{\Sigma}$.
- (iii) ρ maps $\dot{\Sigma}$ to itself and fixes at least one chamber of $\dot{\Sigma}$.

Proof. Let S be as in 5.2. We think of S as a chamber of $\Sigma(\dot{\Phi})$ and let S' be the unique opposite chamber of $\Sigma(\dot{\Phi})$. The map τ fixes both S and S' . Hence if $\dot{\psi}$ is as in (i), then ρ fixes the two opposite chambers $\dot{\psi}(S)$ and $\dot{\psi}(S')$ of $\dot{\Sigma}$. Thus (i) implies (ii). Since opposite chambers are contained in a unique apartment, (ii) implies (iii). Now suppose (iii) holds. Let C be a chamber of $\dot{\Sigma}$ fixed by ρ and let $\dot{\psi}$ be the unique isomorphism from $\Sigma(\dot{\Phi})$ to $\dot{\Sigma}$ mapping S to C . Since the composition $\rho^{-1} \circ \dot{\psi} \circ \tau$ is also a type-preserving automorphism from $\Sigma(\dot{\Phi})$ to $\dot{\Sigma}$ mapping S to C , it must equal $\dot{\psi}$. Thus (i) holds. $\square$

Definition 6.4. Under the hypotheses of 6.3, we call an apartment of $\dot{\Delta}$ ρ -compatible if it satisfies the three equivalent conditions in 6.3.

In the following result, we make implicit use of the fact that a building Δ is completely determined by the graph whose vertices are the chambers of Δ , where two chambers are joined by an edge whenever there is a panel of Δ containing them both. (This is the point of view, for example, in [14].)

Theorem 6.5. *Let $\dot{\Phi}$ and τ be as in 6.1, let Φ be as in 6.2.ii, let $\dot{\Delta}$ be an arbitrary building of type $\dot{\Phi}$ which is not necessarily thick, let ρ be a non-type-preserving automorphism of $\dot{\Delta}$ of order two and let $\dot{\Sigma}$ be an apartment of $\dot{\Delta}$. Suppose that $\dot{\Sigma}$ is ρ -compatible as defined in 6.4. Let $\dot{\psi}$ be as in 6.3 and let π be the bijection from $\dot{\Phi}$ to the set of roots of $\dot{\Sigma}$ induced by the isomorphism $\dot{\psi}$. Let Δ be the graph whose vertices are the chambers of $\dot{\Delta}$ fixed by ρ , where two such chambers are joined by an edge whenever they are opposite in a residue of rank two fixed by ρ and let Σ be the subgraph of Δ spanned by the chambers of $\dot{\Sigma}$ fixed by ρ . Then the following hold:*

- (i) There is a unique isomorphism ψ from $\Sigma(\Phi)$ to Σ such that

$$\psi(\dot{S} \cap \mathbb{A}) = \dot{\psi}(\dot{S})$$

for each τ -invariant sector $\dot{S}$ of $\dot{\Phi}$.

(ii) If π is the bijection from Φ to the set of roots of Σ induced by ψ , then

$$\pi(\ddot{\alpha}/|\ddot{\alpha}|) = \dot{\pi}(\dot{\alpha}) \cap \Sigma$$

for all $\dot{\alpha} \in \dot{\Phi}$, where the map $\dot{\alpha} \mapsto \ddot{\alpha}$ is as in 6.2.

(iii) Δ is a building of type Φ whose apartments are the subgraphs spanned by the chambers fixed by ρ in ρ -compatible apartments of $\dot{\Delta}$.

Proof. Assertions (i) and (ii) hold by 6.2. Next we observe that if R is the ρ -invariant residue of rank two containing two adjacent chambers of Δ , then all the chambers in R fixed by ρ are pairwise opposite in R and hence adjacent in Δ . This means that Δ is a *chamber system* as defined at the beginning of [7]. To show that (iii) holds, it therefore suffices, by Theorem 3.11 in [7] and Theorem 8.21 in [14], to show that every two chambers of Δ are contained in an apartment of $\dot{\Delta}$ containing opposite chambers fixed by ρ . In cases B and G, every two chambers of Δ are opposite in $\dot{\Delta}$. We can thus assume that we are in case F.

Let $\gamma = (C_0, C_1, \dots, C_k)$ be a gallery in Δ . Then for each $i \in [1, k]$, the chambers C_{i-1} and C_i are opposite in a ρ -invariant J_i -residue R_i , where J_i is a two-element subset of the vertex set I of the diagram F_4 invariant under the non-trivial automorphism of this diagram. We will say that γ is *alternating* if $J_i \neq J_{i-1}$ for all $i \in [2, k]$. Let $\dot{\gamma}$ be a gallery in $\dot{\Delta}$ from C_0 to C_k containing C_i also for all $i \in [1, k-1]$ such that the subgallery from C_{i-1} to C_i is a minimal gallery in R_i for each $i \in [1, k]$. Thus the length of $\dot{\gamma}$ is

$$m := d_1 + d_2 + \dots + d_k,$$

where d_i is the diameter of R_i for all $i \in [1, k]$. Now suppose that $k = 8$. Then m equals the diameter of $\dot{\Sigma}$ and if D and D' are opposite chambers in $\dot{\Sigma} \cap \Delta$, then there exists a unique minimal gallery from D to D' that has the same type as $\dot{\gamma}$. By Proposition 7.7.ii in [14], it follows that $\dot{\gamma}$ is minimal and hence C_0 and C_8 are opposite. By Corollaries 8.6 and 8.9 in [14], therefore, $\dot{\gamma}$ is contained in an apartment of $\dot{\Delta}$. It thus suffices to show that every alternating gallery of arbitrary length k in Δ can be extended to an alternating gallery in Δ of length $k+1$ and that every two chambers of Δ are joined by an alternating gallery in Δ .

Suppose first that the gallery $\gamma = (C_0, C_1, \dots, C_k)$ is alternating and let J_i for $i \in [1, k]$ be as in the previous paragraph. Let $J = I \setminus J_k$ and let R be the unique J -residue containing C_k . Since ρ fixes C_k , the residue R is also ρ -invariant. The residue R is a generalized n -gon for $n = 2$ or 4 . Let γ be a gallery in R of length $n/2$ starting at C_k and let γ_1 be the concatenation of γ^{-1} with γ^ρ . Then γ_1 is a minimal galley of length n (because ρ is not type-preserving) and ρ preserves γ_1 (because $\rho^2 = 1$). Therefore ρ fixes the unique chamber C_{k+1} opposite C_k in the unique apartment of R containing γ_1 . Thus $(C_0, \dots, C_k, C_{k+1})$ is an alternating gallery extending γ .

Now suppose that C and C' are two arbitrary chambers of Δ and let e be the distance between them in $\dot{\Delta}$. Since Δ is a chamber system (as observed above), we can obtain an alternating gallery from C to C' from an arbitrary gallery from C to C' simply by discarding superfluous chambers. It will suffice to show, therefore, that there is a gallery from C to C' in Δ . We proceed by induction with respect to e . We can suppose that $e > 0$. Thus we can choose a chamber C_1 adjacent to C' that is at distance $e-1$ from C . Let R be the unique ρ -invariant residue of rank two containing both C and C_1 and let $C_2 = \text{proj}_R C'$. Since ρ fixes C' and

R , it fixes C_2 too. Thus C_2 is a chamber of Δ adjacent to C in Δ and at distance strictly less than e to C' . By induction, we conclude that there is, in fact, a gallery in Δ from C to C' . $\square$

Notation 6.6. Let $\Delta, \Sigma, \psi, \rho$ as well as U_α and x_α be as in 5.3. Note that the type of Δ is, according to 6.1, now $\dot{\Phi}$ rather than Φ . In order to focus on the set Δ^ρ defined in 5.7, we now replace also the designations $\Delta, \Sigma, \psi, U_\alpha$ and x_α by $\dot{\Delta}, \dot{\Sigma}, \dot{\psi}, U_{\dot{\alpha}}$ and $x_{\dot{\alpha}}$ (but let ρ remain ρ). We then set $\Delta = \dot{\Delta}^\rho$ and $\Sigma = \dot{\Sigma} \cap \Delta$. By 6.5 applied to these data, Δ has (canonically) the structure of a building of type Φ , where Φ is as in 6.2.ii, and Σ is an apartment of Δ . Let π be the map obtained from these data in 6.5.ii and for each $\alpha \in \Phi$, let $\dot{\Phi}_\alpha$ denote the pre-image of α under the surjection $\dot{\alpha} \mapsto \ddot{\alpha}/|\ddot{\alpha}|$ from $\dot{\Phi}$ to Φ , where the map $\dot{\alpha} \mapsto \ddot{\alpha}$ is as in 6.2.

Theorem 6.7. *Let $\dot{\Phi}, U_{\dot{\alpha}}$ for $\dot{\alpha} \in \dot{\Phi}, \Phi, \dot{\Phi}_\alpha$ for $\alpha \in \Phi, \Delta, \Sigma, \rho$ and π be as in 6.6 and let Φ be identified with the set of roots of Σ via π . For each $\alpha \in \Phi$, let U_α denote the centralizer of ρ in the subgroup*

$$\langle U_{\dot{\alpha}} \mid \dot{\alpha} \in \dot{\Phi}_\alpha \rangle$$

of $\text{Aut}(\dot{\Delta})$. Then U_α acts faithfully on Δ for each $\alpha \in \Phi$ (in all three cases); $(\Sigma, (U_\alpha)_{\alpha \in \Phi})$ is a Moufang structure on Δ in cases B and G; and in case F, Δ is Moufang and for each $\alpha \in \Phi, U_\alpha$ is the corresponding root group.

Proof. Let $\dot{\Delta}$ be as in 6.6 and let $\alpha \in \Phi$. We think of α as a root of Σ and choose a panel of Σ containing one chamber C in α and another C' not in α . Then C and C' are opposite in a unique rank two residue $\dot{R}$ of $\dot{\Delta}$ fixed by ρ . By 6.5.ii, $\dot{\Phi}_\alpha$ consists of precisely those roots of $\dot{\Sigma}$ that contain C but not C' . By Proposition 8.13 in [14], the map $\dot{\alpha} \mapsto \dot{\alpha} \cap \dot{R}$ is thus a bijection from $\dot{\Phi}_\alpha$ to the set of roots of the apartment $\Sigma \cap \dot{R}$ of $\dot{R}$ and by Proposition 11.10 in [14], $U_{\dot{\alpha}}$ induces the root group on $\dot{R}$ corresponding to the root $\dot{\alpha} \cap \dot{R}$ for each $\dot{\alpha} \in \dot{\Phi}_\alpha$. By Theorem 11.11.ii in [14], therefore, the group

$$\langle U_{\dot{\alpha}} \mid \dot{\alpha} \in \dot{\Phi}_\alpha \rangle$$

acts sharply transitively on the set Q of chambers of $\dot{R}$ which are opposite C in $\dot{R}$. Thus the group U_α acts sharply transitively on the fixed point set Q^ρ of ρ in Q . It follows that U_α acts faithfully on Δ and by Theorem 9.3 in [14], that U_α acts sharply transitively on the set of apartments of Δ containing α . We conclude that $(U_\alpha)_{\alpha \in \Phi}$ is a Moufang structure on Δ (as defined in 3.8) in cases B and G.

Now suppose that we are in case F and choose a panel of Δ containing two chambers C_1 and C'_1 in α . There exists a unique rank two residue $\dot{R}_1$ of $\dot{\Delta}$ fixed by ρ containing C_1 and C'_1 . Let $\dot{P}$ be a panel of $\dot{R}_1$ containing two chambers of $\dot{\Sigma}$. If $\dot{\alpha} \in \dot{\Phi}_\alpha$, then $\dot{\alpha}$ contains both C_1 and C'_1 , hence the apartment $\dot{R}_1 \cap \dot{\Sigma}$ of $\dot{R}_1$ is contained in $\dot{\alpha}$ (since roots are convex) and thus $U_{\dot{\alpha}}$ acts trivially both on $\dot{R}_1 \cap \dot{\Sigma}$ and on $\dot{P}$. By Theorem 9.7 in [14], therefore, $U_{\dot{\alpha}}$ acts trivially on $\dot{R}_1$ for all $\dot{\alpha} \in \dot{\Phi}_\alpha$. It follows that U_α is contained in the root group of Δ corresponding to α . Thus Δ is Moufang since U_α acts transitively (in fact, sharply transitively) on the set of apartments of Δ containing α and α is arbitrary. By Theorem 9.3 and Proposition 11.4 in [14], the root group corresponding to α also acts sharply transitively on the set of apartments of Δ containing α . It follows that U_α equals this root group. $\square$

Convention 6.8. We will sometimes refer to a building Δ of the sort that appears in 6.7 as a *Suzuki-Ree building*. When we say that Δ is a Suzuki-Ree building in cases B or G, we mean that we have in mind the Moufang structure on Δ described in 6.7.

Remark 6.9. Let Δ_F and $\dot{\Delta}_F$ be the buildings called Δ and $\dot{\Delta}$ in 6.6 in case F and let Δ_B and $\dot{\Delta}_B$ denote the buildings called Δ and $\dot{\Delta}$ in 6.6 in case B under the assumption that $L = K$, where L is as in 5.1. Then there exist residues of rank two of $\dot{\Delta}_F$ fixed by ρ which are isomorphic to the building $\dot{\Delta}_B$. Let $\dot{R}$ be one of these residues and let P be the corresponding panel of Δ_F . Then P can be identified with the building Δ_B in such a way that the canonical Moufang structure on P which comes from Δ_F as described in 3.9 coincides with the Moufang structure on Δ_B described in 6.7.

Notation 6.10. Let $S = L \times L$ in case B and let $S = K \times K$ in case F. In both of these two cases, let

$$(s, t) \cdot (u, v) = (s + u, t + v + s^\theta u)$$

and

$$R(s, t) = s^{\theta+2} + st + t^\theta$$

for all $(s, t) \in S$. In case G, let $T = K \times K \times K$, let

$$(r, s, t) \cdot (w, u, v) = (r + w, s + u + r^\theta w, t + v - ru + sw - r^{\theta+1}w)$$

for all $(r, s, t), (w, u, v) \in T$ and let

$$N(r, s, t) = r^{\theta+1}s^\theta - rt^\theta - r^{\theta+3}s - r^2s^2 + s^{\theta+1} + t^2 - r^{2\theta+4}$$

for all $(r, s, t) \in T$. Then S and T are groups (with multiplication $\cdot$),

$$(s, t)^{-1} = (s, t + s^{\theta+1})$$

for all $(s, t) \in S$ and

$$(r, s, t)^{-1} = (-r, -s + r^{\theta+1}, -t)$$

for all $(r, s, t) \in T$. We will call R the *norm of the group S* and N the *norm of the group T* . The center of S is $\{(0, t) \mid t \in K \text{ (or } t \in L)\}$ and the center of T is $\{(0, 0, t) \mid t \in K\}$; both centers are isomorphic to the additive group of K .

(Note that in case B, the product $K^\theta L$ is contained in L (by 5.1) and thus the products $s^\theta t$, $R(s, t)^{2-\theta}u$ and $R(s, t)^\theta v$ are contained in L for all $(s, t), (u, v) \in S$ even though the norm $R(s, t)$ is not necessarily contained in L . See 6.12.i.)

Remark 6.11. It is shown in [9] that the maps R and N are anisotropic. By this we mean that $R(s, t) = 0$ only if $(s, t) = 0$ and $N(r, s, t) = 0$ only if $(r, s, t) = 0$.

Proposition 6.12. Let Δ , Φ , Σ , the identification of Φ with the set of roots of Σ and the root groups U_α be as in 6.7, let G be the corresponding Ree or Suzuki group as defined in 5.7 and let the groups S and T and their norms R and N be as in 6.10. Then there exist $\alpha \in \Phi$ such that the following hold:

- (i) In cases B and F, $U_\alpha \cong S$ and there exists an isomorphism x_α from S to U_α that

$$x_\alpha(u, v)^{h(s, t)} = x_\alpha(R(s, t)^{2-\theta}u, R(s, t)^\theta v)$$

for all $(u, v) \in S$ and all $(s, t) \in S^*$, where

$$h(s, t) = m_\Sigma(x_\alpha(0, 1))m_\Sigma(x_\alpha(s, t)).$$

- (ii) In case G, $U_\alpha \cong T$ and there exists an isomorphism x_α from T to U_α such that

$$x_\alpha(w, u, v)^{h(r, s, t)} = x_\alpha(N(r, s, t)^{2-\theta}w, N(r, s, t)^{\theta-1}u, N(r, s, t)v)$$

for all $(w, u, v) \in T$ and all $(r, s, t) \in T^*$, where

$$h(r, s, t) = m_\Sigma(x_\alpha(0, 0, 1))m_\Sigma(x_\alpha(r, s, t)).$$

The element α is unique up to the action of the stabilizer G_Σ of Σ in G on the set of roots of Σ .

Proof. By [13, 33.17], (i) holds in case F. By 6.9, it follows that (i) holds also in case B when $K = L$. Simply by restricting scalars, we conclude that (i) holds in B also when $L \neq K$.

Suppose now that we are in case G. Let C and C_1 be the two chambers in Σ and suppose that $\alpha = \{C\}$. Let $\dot{\alpha}_i$ be the elements of $\dot{\Phi}$ ordered clockwise modulo 12 so that $\dot{\alpha}_1, \dots, \dot{\alpha}_6$ are the roots in the set $\dot{\Phi}_\alpha$ defined in 6.7 (where $\dot{\Phi}$, $\dot{\alpha}_i$, etc. are as in 6.6). Let U_i denote the root group $U_{\dot{\alpha}_i}$ of $\dot{\Delta}$ for all i , let $(x_i)_{i \in [1, 6]}$ denote the collection of isomorphisms $x_i = x_{\dot{\alpha}_i}$ from the additive group of K to U_i which appear in the relations 5.4–5.6 and let U_+ denote the subgroup of $\text{Aut}(\dot{\Delta})$ generated by the root groups U_i for all $i \in [1, 6]$. By 5.3.i, $x_i(t)^\rho = x_{7-i}(t)$ for all $i \in [1, 6]$ and by 6.7, U_α is the centralizer of ρ in U_+ . Let

$$(6.13) \quad x_\alpha(r, s, t) = x_1(r)x_2(r^{\theta+1} - s)x_3(t + rs)x_4(r^{\theta+2} - rs + t)x_5(-s)x_6(r)$$

for all $(r, s, t) \in T$. Thus x_α is a map from T to U_+ . By 5.3–5.6, 6.10 and a bit of calculation, this map is, in fact, an isomorphism from T to U_α .

Let $G^\dagger$ be the subgroup of $\text{Aut}(\dot{\Delta})$ as defined in 3.11. Since the elements in the stabilizer $G_{C, C_1}^\dagger$ are type-preserving and fix opposite chambers of the apartment $\dot{\Sigma}$, they are contained in the subgroup H of $\text{Aut}(\dot{\Delta})$ defined in 5.13. If $g \in G_{C, C_1}^\dagger$, then by 5.13, there exist $z, z_1, z_2 \in K^*$ such that $x_\alpha(r, s, t)^g = x_\alpha(zr, z_1s, z_2t)$ for all $(r, s, t) \in T$. Since x_α is an isomorphism, the map $(r, s, t) \mapsto (zr, z_1s, z_2t)$ is an automorphism of T . It follows that $z_1 = z^{\theta+1}$ and $z_2 = z^{\theta+2}t$. Thus for each element g of $G^\dagger$ fixing the two chambers C and C_1 of the apartment Σ , there exists an element $z \in K^*$ such that

$$(6.14) \quad x_\alpha(r, s, t)^g = x_\alpha(zr, z^{\theta+1}s, z^{\theta+2}t)$$

for all $(r, s, t) \in T$.

Let the group T be identified with the set $\Delta \setminus \{C\}$ via the map sending $a \in T$ to the image of C_1 under the element $x_\alpha(a)$ of U_α . Even though we are using exponential notation (and, by implication, composition from left to right) in the claim we are proving, for the remainder of this proof we will think of the group G as acting on the set $C \cup T$ from the left (with composition from right to left) in order to conform with the notation in [9] from where we borrow the following argument. For the same reason, we will also use additive notation for T (only in this proof) even though it is not an abelian group; in particular, we let 0 denote the identity $(0, 0, 0)$ of T . Thus $\Sigma = \{C, 0\}$ and for each $a \in T$, the element $x_\alpha(a)$ of U_α fixes C and induces the map $b \mapsto a + b$ on T .

For each element $a = (r, s, t) \in T^*$, we set

$$(6.15) \quad u(a) = r^2s - rt + s^\theta - r^{\theta+3}$$

and

$$(6.16) \quad v(a) = r^\theta s^\theta - t^\theta + rs^2 + st - r^{2\theta+3}.$$

(These expressions are taken from [9, 5.3], where $N(a)$ is called $w = w(a)$. Note that in [9, 5.3], there is an exponent θ missing in the second term of w and a minus sign missing in front of the whole formula for w ; see Section 2.10 in [3].) As is explained in Section 5 of [9], $N(a) \neq 0$ if $a \neq 0$ and there is an element ω in G interchanging the two chambers C and 0 of Σ such that

$$(6.17) \quad \omega(a) = (-v(a)/N(a), -u(a)/N(a), -t/N(a))$$

for all $a = (r, s, t) \in T^*$. By 6.10, 6.15, 6.16 and 6.17 (and a bit of calculation), we have

$$(6.18) \quad N(\omega(a)) = N(a)^{-1}$$

and

$$(6.19) \quad N(-a) = N(a)$$

for all $a \in T^*$ (where $-a$ is the inverse of a in T).

The element ω^2 fixes 0 and C . By 6.14, there thus exists $z \in K^*$ such that

$$(6.20) \quad \omega^2(r, s, t) = (zr, z^{\theta+1}s, z^{\theta+2}t)$$

for all $(r, s, t) \in T$. Let $v = z^{\theta+2}$. Then

$$(0, 0, v) = \omega^2(0, 0, 1) = \omega(1, 0, -1) = (0, 0, 1)$$

by 6.17 and therefore $z = v^{2-\theta} = 1$. We conclude that

$$(6.21) \quad \omega^2 = 1$$

by 6.20.

Now let $a = (r, s, t) \in T^*$ and set $a' = \omega(-\omega(a))$. (This makes sense since ω maps T^* to itself.) By 6.21, the two products $\omega x_\alpha(a)\omega$ and $x_\alpha(\omega(a))\omega x_\alpha(a')$ both map 0 to 0 and C to $\omega(a)$. Thus

$$(6.22) \quad \rho_a := (x_\alpha(\omega(a))\omega x_\alpha(a'))^{-1} \omega x_\alpha(a)\omega$$

fixes both C and 0 . By 6.21 and 6.22, we have

$$(6.23) \quad \rho_a(\omega(-a)) = -a' = -\omega(-\omega(a)).$$

Let $\xi = \rho_a \omega$. Thus ξ is an element of G interchanging C and 0 . By 6.21 and 6.22, we have

$$\xi = x_\alpha(-a') \cdot \omega x_\alpha(-\omega(a))\omega \cdot x_\alpha(a) \in U_\alpha^* \cdot U_{-\alpha}^* \cdot U_\alpha^*.$$

By 3.10.i, therefore,

$$\xi = m_\Sigma(\omega x_\alpha(-\omega(a))\omega)$$

and

$$x_\alpha(a) = \lambda(\omega x_\alpha(-\omega(a))\omega).$$

Hence by 3.10.iii, $\xi = m_\Sigma(x_\alpha(a))$. Therefore

$$(6.24) \quad \rho_a = m_\Sigma(x_\alpha(a))\omega.$$

By 6.14, there exists $z \in K^*$ such that

$$(6.25) \quad \rho_a(w, u, v) = (zw, z^{\theta+1}u, z^{\theta+2}v)$$

for all $(w, u, v) \in T$. By 6.18 and 6.19, we have $N(-\omega(a)) = N(a)^{-1}$. By 6.17 and 6.25, therefore, the third coordinate of $-\omega(-\omega(a))$ is t , whereas the third coordinate of $\rho_a(\omega(-a))$ is $tz^{\theta+2}/N(a)$. By 6.23, it follows that $z^{\theta+2} = N(a)$ and hence

$$(6.26) \quad z = N(a)^{2-\theta}$$

if $t \neq 0$. If $t = 0$, we obtain the same conclusion by comparing the first or second coordinates of both sides of the identity 6.23; we leave these calculations to the reader. By 6.25 and 6.26, finally, we have $\rho_a = 1$ if $a = (0, 0, 1)$. By 6.21 and 6.24, it follows that $\omega = m_\Sigma(0, 0, 1)$. Thus (ii) holds by 6.24, 6.25 and 6.26. (Note that in (ii), $x_\alpha(w, u, v)^{h(r, s, t)}$ is to be interpreted as $x_\alpha(w, u, v)$ conjugated first by $m_\Sigma(x_\alpha(0, 0, 1))$ and then by $m_\Sigma(x_\alpha(r, s, t))$, whereas in the proof $\rho_a(w, u, v)$ is to be interpreted as the image of (w, u, v) under $m_\Sigma(x_\alpha(0, 0, 1))$ to which then the map $m_\Sigma(x_\alpha(r, s, t))$ is applied.) $\square$

7. BRUHAT-TITS SPACES FOR THE REE AND SUZUKI GROUPS

We begin this section with a result which explains why a valuation of the root datum of a Suzuki-Ree building defined over a pair (K, θ) (or triple (K, θ, L)) requires the existence of a θ -invariant valuation of K . We then formulate our most important result in 7.11.

Theorem 7.1. *Let $\Delta, \Sigma, \Phi, \alpha, x_\alpha$, etc. be as in 6.6 and 6.12, let $w = x_\alpha(0, 1)$ in cases B and F and let $w = x_\alpha(0, 0, 1)$ in case G, let ψ be a valuation of the root datum of Δ based at Σ and let*

$$\varphi = \psi - \psi_\alpha(w)\alpha/(\alpha \cdot \alpha)$$

(as defined in 3.15). (Thus φ is a valuation equipollent to ψ such that $\varphi_\alpha(w) = 0$.) Then there exists a unique θ -invariant valuation ν of K , which depends only on the equipollence class of ψ , such that

$$(7.2) \quad \varphi_\alpha(x_\alpha(s, t)) = \nu(R(s, t))$$

for all $(s, t) \in S^*$ in cases B and F and

$$(7.3) \quad \varphi_\alpha(x_\alpha(r, s, t)) = \nu(N(r, s, t))$$

for all $(r, s, t) \in T^*$ in case G.

Proof. Let

$$\nu(t) = \varphi_\alpha(x_\alpha(0, t^\theta))/2$$

for all $t \in K^*$ in cases B and F and let

$$\nu(t) = \varphi_\alpha(x_\alpha(0, 0, t))/2$$

for all $t \in K^*$ in case G. Then 7.2 and 7.3 hold by 3.18 with $g = w$ and $u = x_\alpha(s, t)$ or $x_\alpha(r, s, t)$ and 6.12. It thus need only to show that ν is a θ -invariant valuation of K .

Let $\nu(0) = \infty$. By (V1), we have $\nu(s + t) \geq \min\{\nu(s), \nu(t)\}$ for all $s, t \in K$. By 6.12 again, we have

$$x_\alpha(0, s^\theta)^{m_\Sigma(w)m_\Sigma(x_\alpha(0, t^\theta))} = x_\alpha(0, s^\theta t^{2\theta})$$

for all $s, t \in K^*$ in cases B and F and

$$x_\alpha(0, 0, s)^{m_\Sigma(w)m_\Sigma(x_\alpha(0, 0, t))} = x_\alpha(0, 0, st^2)$$

for all $s, t \in K^*$ in case G. By 3.18 again, this time with $g = x_\alpha(0, t^\theta)$ or $x_\alpha(0, 0, t)$ and $u = x_\alpha(0, s^\theta)$ or $x_\alpha(0, 0, s)$, it follows that

$$\nu(s^2 t) = \nu(t) + 2\nu(s)$$

for all $s, t \in K^*$ in all three cases. From this identity, we thus obtain

$$\nu(s^2 t^2) = \nu(t^2) + 2\nu(s)$$

for all $s, t \in K^*$ and (setting $t = 1$)

$$\nu(s^2) = 2\nu(s)$$

for all $s \in K^*$. Therefore

$$\nu(st) = \nu(s) + \nu(t)$$

for all $s, t \in K^*$. Since $|\varphi_\alpha(U_\alpha^*)| > 1$ by 3.14, it follows from 7.2 and 7.3 that $|\nu(K^*)| > 1$. Thus ν is a valuation of K .

It remains only to show that $\nu(u) \geq 0$ implies that $\nu(u^\theta) \geq 0$. Let u be an element of K^* such that $\nu(u) \geq 0$ (and hence $\nu(u^2) = 2\nu(u) \geq 0$). Suppose first that we are in case B or F, so

$$(1, 0) \cdot (0, u^\theta) = (1, u^\theta)$$

in S . By (V1), therefore,

$$\varphi_\alpha(x_\alpha(1, u^\theta)) \geq \min\{\varphi_\alpha(x_\alpha(1, 0)), \varphi_\alpha(x_\alpha(0, u^\theta))\}.$$

Hence by 7.2,

$$\nu(1 + u^\theta + u^2) \geq \min\{0, \nu(u^2)\} = 0.$$

It follows that $\nu(u^\theta) \geq 0$.

Suppose now that we are in case G, so

$$(1, 0, 0) \cdot (0, 0, u) = (1, 0, u)$$

in T . By (V1), therefore,

$$\varphi_\alpha(x_\alpha(1, 0, u)) \geq \min\{\varphi_\alpha(x_\alpha(1, 0, 0)), \varphi_\alpha(x_\alpha(0, 0, u))\}.$$

Hence by 7.3,

$$\nu(u^2 - u^\theta - 1) \geq \min\{0, \nu(u^2)\} = 0.$$

Again we conclude that $\nu(u^\theta) \geq 0$. $\square$

The converse of 7.1 is also valid:

Theorem 7.4. *Let $\Delta, \Sigma, \Phi, \alpha, x_\alpha$, etc. be as in 6.6 and 6.12. Suppose that ν is a θ -invariant valuation of K and let φ_α be the map given in 7.2 (in cases B and F) or 7.3 (in case G). Then φ_α extends to a valuation of the root datum of Δ based at Σ .*

It would not be hard to prove this result directly. The principal difficulty is to show that

$$(7.5) \quad \nu(R((s, t) \cdot (u, v))) \geq \min\{\nu(R(s, t)), \nu(R(u, v))\}$$

for all $(s, t), (u, v) \in S$ and

$$(7.6) \quad \nu(N((r, s, t) \cdot (w, u, v))) \geq \min\{\nu(N(r, s, t)), \nu(N(w, u, v))\}$$

for all $(r, s, t), (w, u, v) \in T$. These inequalities are required to verify (V1).

Rather than prove 7.4 directly, however, we will prove a stronger result (7.7–7.11) which will have 7.4 (and thus also the two inequalities 7.5 and 7.6) as corollaries.

In Section 9 we include a direct proof of the inequalities 7.5 and 7.6 only because it might be of some independent interest. See also 9.1.10 in [2].

Notation 7.7. Let $\dot{\Phi}$, $\dot{\mathbb{A}}$, $\dot{\mathbb{A}}$ and τ be as in 6.1, let $\dot{\Delta}$, Δ , $\dot{\Sigma}$ and Σ be as in 6.6, let $\dot{W}$ be the Weyl group of $\dot{\Phi}$, let W be the restriction of the centralizer of τ in $\dot{W}$ to the subspace $\mathbb{A}$ and let $\mathbb{W}$ be the group of all isometries of $\mathbb{A}$ generated by W and all translations of $\mathbb{A}$. Let ν be a θ -invariant valuation of K , let ρ be the automorphism of $\dot{\Delta}$ described in 5.3.i, let $\dot{\Phi}$ be identified with the set of roots of $\dot{\Sigma}$ via the map called $\dot{\pi}$ in 6.5 and let $\dot{\varphi}$ denote the ρ -invariant valuation of the root datum of $\dot{\Delta}$ based at $\dot{\Sigma}$ determined by ν as described in 5.15. Let $(\dot{X}, \dot{\mathcal{A}})$ be the Bruhat-Tits space of type $\dot{\Phi}$, $\dot{A}$ the apartment of $(\dot{X}, \dot{\mathcal{A}})$ and x_A the point of $\dot{A}$ obtained by applying 4.16 to $\dot{\Delta}$, $\dot{\Sigma}$ and $\dot{\varphi}$. The pair $(\dot{X}, \dot{\mathcal{A}}_\tau)$ defined as in 4.4 can also be thought of as the Bruhat-Tits space of type $\dot{\Phi}$ obtained by applying 4.16 to $\dot{\Delta}$, $\dot{\Sigma}$ and $\dot{\varphi}$, but only after identifying $\dot{\Phi}$ with the set of roots of $\dot{\Sigma}$ via $\dot{\pi} \circ \tau$ rather than $\dot{\pi}$. By the uniqueness assertion in 4.16, there exists a τ -automorphism $\dot{\rho}$ of $(\dot{X}, \dot{\mathcal{A}})$ (as defined in 4.4) that induces the automorphism ρ on $\dot{\Delta}$ and maps the pair $(\dot{A}, x_A)$ to itself. This map satisfies

$$(7.8) \quad \dot{\rho} \circ \dot{f}_{\dot{A}, x_A} = \dot{f}_{\dot{A}, x_A} \circ \tau,$$

where $\dot{f}_{\dot{A}, x_A} \in \dot{\mathcal{A}}$ is as defined in 4.8. Let $\dot{\mathcal{A}}_\rho$ denote the set of charts $\dot{f} \in \dot{\mathcal{A}}$ such that $\dot{\rho}$ maps the apartment $\dot{f}(\dot{\mathbb{A}})$ to itself and acts trivially on $\dot{f}(\dot{\mathbb{A}})$, let

$$\mathcal{A} = \{\dot{f}|_{\mathbb{A}} \mid \dot{f} \in \dot{\mathcal{A}}_\rho\},$$

and let

$$(7.9) \quad X = \bigcup_{\dot{f} \in \dot{\mathcal{A}}_\rho} \dot{f}(\dot{\mathbb{A}}).$$

Thus X is contained in the fixed point set $\dot{X}^{\dot{\rho}}$ of $\dot{\rho}$. By 7.8, $\dot{f}_{\dot{A}, x_A} \in \dot{\mathcal{A}}_\rho$ and hence

$$(7.10) \quad A := \dot{f}_{\dot{A}, x_A}(\dot{\mathbb{A}})$$

is a subset of X and $x_A = \dot{f}_{\dot{A}, x_A}(0)$ is a point of A .

Here now is our main result:

Theorem 7.11. *Let Φ , ν , $(X, \mathcal{A})$, A and x_A be as in 7.7. Thus, in particular, ν is a θ -invariant valuation of K . Then the following hold:*

- (i) *The pair $(X, \mathcal{A})$ is a Bruhat-Tits space of type Φ whose building at infinity is Δ (in the sense of 4.14 if the rank of Δ is one) and A is an apartment of $(X, \mathcal{A})$.*
- (ii) *Let $\alpha \in \Phi$ be as in 6.12 and let φ be the valuation of the root datum of Δ based at Σ that appears in 4.12.ii when 4.12 (and then 4.15) is applied to the triple $(X, \mathcal{A})$, A and x_A . Then there exists a valuation ν_1 equivalent to ν such that φ_α satisfies 7.2 or 7.3 with ν_1 in place of ν .*

Note that 7.4 is a consequence of 7.11.

8. THE PROOF OF 7.11

For the rest of this paper, we let $\dot{\Delta}$, Δ , $(\dot{X}, \dot{\mathcal{A}})$, $\dot{\Phi}$, $(\dot{\mathbb{A}}, \dot{\mathbb{W}})$, τ , ρ , Φ , $(\mathbb{A}, \mathbb{W})$, $(X, \mathcal{A})$, A , x_A , $\dot{\rho}$, $\dot{f}_\rho$, etc. be as in 7.7. Let $\dot{X}^{\dot{\rho}}$ denote the set of fixed points of $\dot{\rho}$. We prove 7.11 in a series of steps.

Proposition 8.1. *Suppose that $\dot{A}_1$ is an apartment of $(\dot{X}, \dot{A})$. Then the following are equivalent:*

- (i) $\dot{A}_1$ is $\dot{\rho}$ -invariant and contains at least one sector fixed by $\dot{\rho}$.
- (ii) There exists a chart $\dot{f}_1$ in $\dot{A}_\rho$ such that $\dot{\rho} \circ \dot{f}_1 = \dot{f}_1 \circ \tau$.
- (iii) $\dot{A}_1$ is the image of a chart in $\dot{A}_\rho$.

If $\dot{f}_1$ is as in (ii), then $\dot{X}^{\dot{\rho}} \cap \dot{A}_1 = X \cap \dot{A}_1 = \dot{f}_1(\mathbb{A})$.

Proof. Let $\dot{f}$ be a chart in $\dot{A}$ such that $\dot{A}_1 = \dot{f}(\dot{\mathbb{A}})$. Suppose first that $\dot{A}_1$ is $\dot{\rho}$ -invariant and that there exists a sector S of $\dot{\Phi}$ such that

$$(8.2) \quad \dot{\rho}(\dot{f}(S)) = \dot{f}(S).$$

By 4.4, there exists $\dot{f}' \in \dot{A}$ such that $\dot{\rho} \circ \dot{f} = \dot{f}' \circ \tau$. Since $\dot{A}_1$ is $\dot{\rho}$ -invariant, also $\dot{f}'(\dot{\mathbb{A}})$ equals $\dot{A}_1$. By (A2) (in 4.2), therefore, there exists $\dot{w} \in \dot{\mathbb{W}}$ such that $\dot{f}' = \dot{f} \circ \dot{w}$. Thus

$$(8.3) \quad \dot{\rho} \circ \dot{f} = \dot{f} \circ \dot{w} \circ \tau.$$

By 8.2, it follows that $\dot{w} \circ \tau$ fixes S . Thus $\dot{w} \circ \tau$ is a non-trivial automorphism of $\Sigma(\dot{\Phi})$ fixing S . There is only one such automorphism. Since τ also fixes a sector of $\dot{\Phi}$ and $\dot{W}$ acts transitively on the set of sectors of $\dot{\Phi}$, it follows that there exists $\dot{w}_1 \in \dot{W}$ such that $\dot{w} \circ \tau = \dot{w}_1 \circ \tau \circ \dot{w}_1^{-1}$. Let $\dot{f}_1 = \dot{f} \circ \dot{w}_1$. Then $\dot{f}_1 \in \dot{A}$ by (A1) and

$$\begin{aligned} \dot{\rho} \circ \dot{f}_1 &= \dot{\rho} \circ \dot{f} \circ \dot{w}_1 \\ &= \dot{f} \circ \dot{w} \circ \tau \circ \dot{w}_1 \\ &= \dot{f} \circ \dot{w}_1 \circ \tau = \dot{f}_1 \circ \tau \end{aligned}$$

by 8.3. It follows from this identity that $\dot{\rho}$ acts trivially on $\dot{f}_1(\mathbb{A})$, so $\dot{f}_1 \in \dot{A}_\rho$ (but $\dot{\rho}$ does not fix any other points in $\dot{A}_1$, so $\dot{X}^{\dot{\rho}} \cap \dot{A}_1 = X \cap \dot{A}_1 = \dot{f}_1(\mathbb{A})$). Thus (i) implies (ii). It now suffices to observe that if $\dot{f}_1$ is a chart in $\dot{A}_\rho$ whose image is $\dot{A}_1$, then by 6.2.iv, $\dot{\rho}$ fixes $\dot{f}_1(S)$ for every τ -invariant sector S of $\dot{\Phi}$. $\square$

Proposition 8.4. *Let $\dot{f}$ and $\dot{f}_1$ be two charts in $\dot{A}_\rho$ such that $\dot{A}_1 := \dot{f}(\dot{\mathbb{A}}) = \dot{f}_1(\dot{\mathbb{A}})$. Then*

$$\dot{f}(\mathbb{A}) = \dot{f}_1(\mathbb{A}) = \dot{X}^{\dot{\rho}} \cap \dot{A}_1 = X \cap \dot{A}_1.$$

Proof. By 8.1, it suffices to assume that $\dot{\rho} \circ \dot{f}_1 = \dot{f}_1 \circ \tau$ and $\dot{X}^{\dot{\rho}} \cap \dot{A}_1 = X \cap \dot{A}_1 = \dot{f}_1(\mathbb{A})$. Thus $\dot{f}(\mathbb{A}) \subset \dot{X}^{\dot{\rho}} \cap \dot{A}_1 = \dot{f}_1(\mathbb{A})$. By (A2), therefore, there exists $\dot{w} \in \dot{\mathbb{W}}$ such that $\dot{f}_1 \circ \dot{w}$ and $\dot{f}$ coincide on $\mathbb{A}$. Thus $\dot{f}_1(\dot{w}(\mathbb{A})) = \dot{f}(\mathbb{A}) \subset \dot{f}_1(\mathbb{A})$ and hence $\dot{w}$ maps $\mathbb{A}$ to itself. Therefore $\dot{f}_1(\mathbb{A}) = \dot{f}(\mathbb{A})$. $\square$

Proposition 8.5. *Let $x \in X$ and let g_x be as in 4.10. Then the set of fixed points of $\dot{\rho}$ in the residue $(\dot{X}, \dot{A})_x$ has the structure of a building of type Φ and for all $\dot{f} \in \dot{A}_\rho$ mapping 0 to x , $g_x(\dot{f}(\mathbb{A}))$ is an apartment of this building.*

Proof. By 4.10, $(\dot{X}, \dot{A})_x$ is a building of type Φ whose apartments are all of the form $g_x(\dot{f}(\dot{\mathbb{A}}))$ for some $\dot{f} \in \dot{A}$. The conclusion holds, therefore, by applying 6.5 to this building and the automorphism of this building induced by $\dot{\rho}$. $\square$

Corollary 8.6. *Let $x \in X$, let $\dot{u}$ be a germ at x fixed by $\dot{\rho}$ and let $\dot{f}$ be a chart in $\dot{A}_\rho$ mapping 0 to x . Then there exists a sector with vertex x in the apartment $g_x(\dot{f}(\dot{\mathbb{A}}))$ that is fixed by $\dot{\rho}$ and whose germ is opposite a maximal germ at x containing $\dot{u}$.*

Proof. The germ $\dot{u}$ is a face and $g_x(\dot{f}(\mathcal{A}))$ is an apartment of the building of type Φ described in 8.5. Since the apartment $g_x(\dot{f}(\dot{\mathbb{A}}))$ is $\dot{\rho}$ -invariant, a sector with vertex x in this apartment is fixed by $\dot{\rho}$ if and only if its germ is fixed by $\dot{\rho}$. The claim follows, therefore, from the fact that for each chamber C and each apartment Σ in a spherical building, there always exists a chamber in Σ which is opposite C . $\square$

Proposition 8.7. *Let $\dot{A}_1$ be the image of a chart $\dot{f}_1$ in $\dot{\mathcal{A}}_\rho$. Then the map*

$$\dot{S}_1 \mapsto \dot{S}_1 \cap X$$

is a bijection from the set of sectors of $\dot{A}_1$ that are fixed by $\dot{\rho}$ to the set of sectors of $\dot{f}_1(\dot{\mathbb{A}})$, i.e. to the set of images under $\dot{f}_1$ of sectors of Φ .

Proof. By 6.2, the map $\dot{S} \mapsto \dot{S} \cap \mathbb{A}$ is a bijection from the set of sectors of $\dot{\Phi}$ that are fixed by τ to the set of sectors of Φ . If $\dot{S}$ is any one of these sectors, then $\dot{f}_1(\dot{S} \cap \mathbb{A}) = \dot{f}_1(\dot{S}) \cap X$ by 8.4. $\square$

Proposition 8.8. *The pair $(X, \mathcal{A})$ is a non-discrete Euclidean building of type Φ .*

Proof. We need to show that $(X, \mathcal{A})$ satisfies the conditions (A1)–(A6) formulated in 4.2.

Let $f \in \mathcal{A}$ and $w \in \mathbb{W}$. There exist elements $\dot{f}$ in $\dot{\mathcal{A}}_\rho$ and $\dot{w}$ in the centralizer of τ in $\dot{\mathbb{W}}$ such that f is the restriction of $\dot{f}$ to $\mathbb{A}$ and w is the restriction of $\dot{w}$ to $\mathbb{A}$. Since $(\dot{X}, \dot{\mathcal{A}})$ satisfies (A1), we have $\dot{f} \circ \dot{w} \in \dot{\mathcal{A}}$. It follows that $f \circ w \in \dot{\mathcal{A}}_\rho$ since $\dot{f}(\dot{w}(\dot{\mathbb{A}})) = \dot{f}(\dot{\mathbb{A}})$ and $\dot{f}(\dot{w}(\mathbb{A})) = \dot{f}(\mathbb{A})$. Thus $(X, \mathcal{A})$ satisfies (A1).

Next let $\dot{f}, \dot{f}' \in \dot{\mathcal{A}}_\rho$. Since $(\dot{X}, \dot{\mathcal{A}})$ satisfies (A2), the set

$$\dot{M} := \{v \in \dot{\mathbb{A}} \mid \dot{f}(v) \in \dot{f}'(\dot{\mathbb{A}})\}$$

is closed and convex and there exists $\dot{w} \in \dot{\mathbb{W}}$ such that the maps $\dot{f}$ and $\dot{f}' \circ \dot{w}$ coincide on $\dot{M}$. Let

$$M := \{v \in \mathbb{A} \mid \dot{f}(v) \in \dot{f}'(\mathbb{A})\}.$$

Then $M \subset \dot{M} \cap \mathbb{A}$. Let $v \in \dot{M} \cap \mathbb{A}$. Thus $v \in \mathbb{A}$ and $\dot{f}(v) = \dot{f}'(v')$ for some $v' \in \dot{\mathbb{A}}$. Since $\dot{f} \in \dot{\mathcal{A}}_\rho$, the point $\dot{f}(v)$ is a point of $\dot{f}'(\dot{\mathbb{A}})$ fixed by $\dot{\rho}$. Since also $\dot{f}' \in \dot{\mathcal{A}}_\rho$, it follows by 8.4 that $v' \in \mathbb{A}$ and hence $v \in M$. We conclude that $M = \dot{M} \cap \mathbb{A}$. Since $\dot{M}$ is closed and convex, it follows that M is also closed and convex.

To finish showing that $(X, \mathcal{A})$ satisfies (A2), we can assume that $|M| > 1$. By (A1), we can assume further that the origin 0 is contained in M and that $\dot{w}$ fixes 0, so $\dot{w} \in \dot{W}$. Let $\dot{f}_*$ and $\dot{f}'_*$ be as in 4.11 and let Ξ denote the building of type Φ described in 8.5. By 6.2.iv, we can think of the τ -invariant faces of $\Sigma(\Phi)$ as the faces of $\Sigma(\Phi)$. Thus $\dot{f}_*$ and $\dot{f}'_*$ both map $\Sigma(\Phi)$ to apartments of the building Ξ . In cases B and G, an apartment of Ξ is an arbitrary two-element set of chambers. In case F, an apartment of Ξ is a circuit consisting of 16 chambers and 16 panels and two distinct apartments intersect either in a connected piece of this circuit (possibly empty) or in two opposite panels. Let Y be the set of all faces of $\Sigma(\Phi)$ which are mapped by $\dot{f}_*$ to $\dot{f}'_*(\Sigma(\Phi))$. Thus either $Y = \Sigma(\Phi)$, Y is a simplicial arc in $\Sigma(\Phi)$ (possibly empty) or we are in case F and Y consists of two opposite panels (i.e. two opposite non-maximal faces). The map $(\dot{f}'_*)^{-1} \circ \dot{f}_*$ from Y into $\Sigma(\Phi)$ is type-preserving (since the maps $\dot{f}_*$ and $\dot{f}'_*$ are type-preserving) and if Y consists of two opposite panels of a given type, then $(\dot{f}'_*)^{-1}(\dot{f}_*(Y))$ consists of two opposite panels of the same type (since the maps $\dot{f}_*$ and $\dot{f}'_*$ both map opposite panels to

opposite panels). Thus in every case there exists an element $\dot{w}_1$ in the centralizer of τ in $\dot{W}$ such that $\dot{f}_*$ and $\dot{f}'_* \circ \dot{w}_1$ coincide on Y .

Let z be an arbitrary non-zero element of M and let $z' = \dot{w}(z)$. Since M is convex, it contains the closed interval $[0, z]$. Let $\dot{F}$ be the unique minimal face of $\Sigma(\Phi)$ that contains $[0, z]$. Since $\dot{f}$ and $\dot{f}' \circ \dot{w}$ coincide on M , we have $z' \in \mathbb{A}$ and $\dot{f}(z) = \dot{f}'(z')$. If $\dot{F}'$ is the minimal face of $\Sigma(\Phi)$ that contains $[0, z']$, then $\dot{f}(\dot{F}) = \dot{f}'(\dot{F}')$. Therefore $\dot{F} \in Y$ (and, in particular, Y is not empty). Thus $\dot{f}'(\dot{F}') = \dot{f}_*(\dot{F}) = \dot{f}'_*(\dot{w}_1(\dot{F}))$ by the conclusion of the previous paragraph and hence $\dot{F}' = \dot{w}_1(\dot{F})$. It follows that $\dot{w}(z) = z' \in \dot{w}_1(\dot{F})$, so the points z and $\dot{w}_1^{-1}\dot{w}(z)$ are both contained in the face $\dot{F}$. Since every point of $\mathbb{A}$ distinct from the origin is contained in at most one face of $\dot{\Phi}$ of a given type and the stabilizer of a face in $\dot{W}$ acts trivially on that face, it follows that $\dot{w}_1(z) = \dot{w}(z)$. We conclude that $\dot{f}' \circ \dot{w}_1$ coincides with $\dot{f}$ on M . Thus $(X, \mathcal{A})$ satisfies (A2).

Now let $x, x' \in X$. Since $(\dot{X}, \dot{\mathcal{A}})$ satisfies (A3), there exists an apartment $\dot{A}_1$ of $(\dot{X}, \dot{\mathcal{A}})$ containing the interval $[x, x']$ (as defined in 4.6). By 7.9, there exist $\dot{f}, \dot{f}' \in \dot{\mathcal{A}}_\rho$ such that $x \in \dot{f}(\mathbb{A})$ and $x' \in \dot{f}'(\mathbb{A})$. Since ρ fixes x and x' , it fixes the point $g_x(x')$ of the residue $(\dot{X}, \dot{\mathcal{A}})_x$ defined in 4.10. Let $\dot{F}$ be the unique minimal face of the apartment $\dot{A}_1$ with vertex x whose germ contains the point $g_x(x')$ and let $\dot{u}$ denote the germ of $\dot{F}$. The germ $\dot{u}$ is fixed by ρ (since otherwise $\dot{u}^\rho$ would be disjoint from $\dot{u}$). By 8.6, the apartment $g_x(\dot{f}(\mathbb{A}))$ contains a sector with vertex x which is both fixed by ρ and whose germ is opposite a maximal germ at x containing $\dot{u}$. By Lemma 1.13 in [6], there exists an apartment $\dot{A}_2$ containing $\dot{S}_1 \cup \dot{F}$. Thus $[x, x'] \subset \dot{F} \subset \dot{A}_2$. The convex closure of $\dot{S}_1 \cup \{x'\}$ is a sector of $\dot{A}_2$ with vertex x' which contains the interval $[x, x']$. We denote this sector by $\dot{S}_2$. Since ρ fixes $\dot{S}_1$ and x' , it fixes $\dot{S}_2$ as well. By a second application of 8.6, there exists a sector $\dot{S}_3$ with vertex x' in the apartment $\dot{f}'(\mathbb{A})$ that is fixed by ρ and whose germ is opposite the germ of $\dot{S}_2$ at x' . By Lemma 1.12 in [6], there exists a unique apartment $\dot{A}_3$ containing $\dot{S}_2 \cup \dot{S}_3$. This apartment contains $[x, x']$ (since $\dot{S}_2$ contains $[x, x']$) and is ρ -invariant (since $\dot{S}_2$ and $\dot{S}_3$ are ρ -invariant). By 8.1, there exists a chart $\dot{f}_1$ in $\dot{\mathcal{A}}_\rho$ such that $\dot{A}_3 = \dot{f}_1(\mathbb{A})$ and $x, x' \in \dot{f}_1(\mathbb{A})$. Thus $(X, \mathcal{A})$ satisfies (A3).

Let S and S' be two sectors of X . By 8.7, there exist ρ -invariant sectors $\dot{S}$ and $\dot{S}'$ of $(\dot{X}, \dot{\mathcal{A}})$ such that $S = \dot{S} \cap X$ and $S' = \dot{S}' \cap X$. The chambers $\dot{S}^\infty$ and $(\dot{S}')^\infty$ of $(\dot{X}, \dot{\mathcal{A}})^\infty$ are contained in a ρ -invariant apartment of $(\dot{X}, \dot{\mathcal{A}})^\infty$. This apartment is of the form $\dot{A}_1^\infty$, where $\dot{A}_1$ is a ρ -invariant apartment of $(\dot{X}, \dot{\mathcal{A}})$. The apartment $\dot{A}_1$ contains sectors $\dot{S}_1$ and $\dot{S}'_1$ such that $\dot{S}_1 \subset \dot{S}$ and $\dot{S}'_1 \subset \dot{S}'$. Let $\dot{S}_2 = \dot{S}_1 \cap \rho(\dot{S}_1)$ and $\dot{S}'_2 = \dot{S}'_1 \cap \rho(\dot{S}'_1)$. The intersection of two subsectors of a given sector is again a subsector. It follows that $\dot{S}_2$ is a subsector of $\dot{S}$ and $\dot{S}'_2$ is a subsector of $\dot{S}'$. Since $\rho^2 = 1$, both of these subsectors are fixed by ρ . By 8.1, therefore, $\dot{A}_1 = \dot{f}_1(\mathbb{A})$ for some $\dot{f}_1 \in \dot{\mathcal{A}}_\rho$. Hence by 8.7, $\dot{S}_2 \cap X$ and $\dot{S}'_2 \cap X$ are subsectors of S and S' both are contained in $\dot{f}_1(\mathbb{A})$. Thus $(X, \mathcal{A})$ satisfies (A4).

We turn now to (A5). A root of $(X, \mathcal{A})$ is the image under a chart in $\dot{\mathcal{A}}_\rho$ of H_α for some $\alpha \in \Phi$, where H_α is as in 3.4. Let $\dot{A}_1$ and $\dot{A}_2$ be the images of two charts in $\dot{\mathcal{A}}_\rho$ such that $\dot{A}_1 \cap \dot{A}_2 \cap X$ contains a root β of $(X, \mathcal{A})$ but $\dot{A}_1 \cap X \neq \dot{A}_2 \cap X$. By (A2), the set $\dot{A}_1 \cap \dot{A}_2 \cap X$ is closed. We can therefore choose a point x in this set such that the apartments $\dot{\Sigma}_1 := g_x(\dot{A}_1)$ and $\dot{\Sigma}_2 := g_x(\dot{A}_2)$ of the residue of $(\dot{X}, \dot{\mathcal{A}})_x$ are distinct. For $i = 1$ and 2 , let $\dot{\Sigma}_i^\rho$ be the set of chambers of $\dot{\Sigma}_i$ fixed by ρ . Then

$\dot{\Sigma}_1^\rho$ and $\dot{\Sigma}_2^\rho$ both span apartments of the building Ξ of type Φ defined in 8.5. Let S be a sector contained in the root β . Then the convex hull S' of $\{x\} \cup S$ is a sector with vertex x and $\dot{\Sigma}_1^\rho$ and $\dot{\Sigma}_2^\rho$ both contain the unique chamber of Ξ that contains $g_x(S')$. It follows that $\dot{\Sigma}_1^\rho \cap \dot{\Sigma}_2^\rho$ contains a root of both $\dot{\Sigma}_1^\rho$ and $\dot{\Sigma}_2^\rho$. On the other hand, $\dot{\Sigma}_1^\rho \neq \dot{\Sigma}_2^\rho$ since otherwise $\dot{\Sigma}_1 \cap \dot{\Sigma}_2$ would contain a pair of opposite chambers. Two distinct apartments of a spherical building whose intersection contains a root intersect in a root and their symmetric difference spans a third apartment. Thus the symmetric difference of $\dot{\Sigma}_1^\rho$ and $\dot{\Sigma}_2^\rho$ spans a third apartment of Ξ . Let $\dot{u}_1$ and $\dot{u}_2$ be two chambers in this apartment that are opposite in Ξ and hence also opposite in the residue $(X, \mathcal{A})_x$. There exist unique sectors $\dot{S}_1$ and $\dot{S}_2$ of $\dot{A}_1$ and $\dot{A}_2$ with vertex x whose germs are $\dot{u}_1$ and $\dot{u}_2$, and by Proposition 1.12 in [6] there exists an apartment $\dot{A}_3$ containing these two sectors. In particular, $x \in \dot{A}_1 \cap \dot{A}_2 \cap \dot{A}_3 \cap X$. Now suppose that $\dot{A}'$ is the image of a chart $\dot{f}'$ in $\dot{\mathcal{A}}_\rho$ such that both $\dot{A}' \cap \dot{A}_1 \cap X$ and $\dot{A}' \cap \dot{A}_2 \cap X$ contain roots of $(X, \mathcal{A})$ that are disjoint from $\dot{A}_1 \cap \dot{A}_2 \cap X$. By 8.7, $\dot{f}'(\mathbb{A})$ contains subsectors of both $\dot{S}_1 \cap X$ and $\dot{S}_2 \cap X$ and hence $\dot{A}'$ contains subsectors of both $\dot{S}_1$ and $\dot{S}_2$. Thus $\dot{A}' = \dot{A}_3$ since $\dot{A}_3$ is the convex hull of any two sectors, one contained in S_1 and the other in S_2 . Therefore $x \in \dot{A}_1 \cap \dot{A}_2 \cap \dot{A}'$. Thus $(X, \mathcal{A})$ satisfies (A5).

Let $\dot{d}$ be the metric on $\dot{X}$ that appears in (A6). The restriction of $\dot{d}$ to X is a metric on X , and each chart in $\mathcal{A}$ is the restriction to $\mathbb{A}$ of a chart in $\dot{\mathcal{A}}$. Therefore $(X, \mathcal{A})$ satisfies (A6) with the restriction of $\dot{d}$ to X in place of d . $\square$

Proposition 8.9. *The building Δ is the building at infinity of $(X, \mathcal{A})$ (in the sense of 4.14 in cases B and G).*

Proof. By 8.7, there is a canonical bijection π from the chamber set of $(X, \mathcal{A})^\infty$ to the set of all chambers $\dot{S}^\infty$ of $(\dot{X}, \dot{\mathcal{A}})^\infty = \dot{\Delta}$ such that $\dot{S}$ is a sector of $(\dot{X}, \dot{\mathcal{A}})$ that is fixed by $\dot{\rho}$. If $\dot{S}_1$ is an arbitrary sector of $(\dot{X}, \dot{\mathcal{A}})$ such that $\dot{S}_1^\infty$ is fixed by $\dot{\rho}$, then $\dot{S}_2 := \dot{S}_1 \cap \dot{S}_1^\rho$ is a sector of $(\dot{X}, \dot{\mathcal{A}})$ fixed by $\dot{\rho}$ such that $\dot{S}_2^\infty = \dot{S}_1^\infty$. It follows that π is, in fact, an isomorphism from $(X, \mathcal{A})^\infty$ to Δ . Furthermore, the conclusions of 4.12 hold for $(X, \mathcal{A})$ and Δ , even in cases B and G, by 4.12 applied to $(\dot{X}, \dot{\mathcal{A}})$ and 6.7. $\square$

By 8.8 and 8.9, we conclude that 7.11.i holds. Now let $\alpha \in \Phi$ and φ be as in 7.11.ii.

Proposition 8.10. *There exists a positive real number k (which depends on the case) such that*

$$\varphi_\alpha(x_\alpha(0, t)) = k \cdot \nu(t)$$

for all $t \in K^*$ (or $t \in L^*$) in case B or F and

$$\varphi_\alpha(x_\alpha(0, 0, t)) = k \cdot \nu(t)$$

for all $t \in K^*$ in case G.

Proof. Suppose first that we are in case G. We use the notation from the proof of 6.12. By 6.13, in particular, we have

$$x_\alpha(0, 0, t) = x_{\dot{\alpha}_4}(t)x_{\dot{\alpha}_5}(t)$$

for all $t \in K$. Since τ interchanges $\dot{\alpha}_4$ and $\dot{\alpha}_5$, there exists a positive real number k such that for all $t \in K^*$, the affine half-spaces $H_{\dot{\alpha}_4, \nu(t)}$ and $H_{\dot{\alpha}_5, \nu(t)}$ (as defined in

4.12) both intersect $\mathbb{A}$ (which is the space of fixed points of τ) in an affine half-space of the form $H_{\alpha, k \cdot \nu(t)}$. Choose $t \in K^*$, let $u = x_{\dot{\alpha}_4}(t)$, let $y = x_{\dot{\alpha}_5}(t)$ and let $z = uy$. Then

$$\dot{A} \cap \dot{A}^u = \dot{f}_{\dot{A}, x_A}(H_{\dot{\alpha}_4, \nu(t)})$$

and

$$\dot{A} \cap \dot{A}^y = \dot{f}_{\dot{A}, x_A}(H_{\dot{\alpha}_5, \nu(t)})$$

by 4.12.ii and 5.15. Suppose that z fixes a point x in $\mathbb{A}$ which is not in

$$\dot{f}_{\dot{A}, x_A}(H_{\alpha, k \cdot \nu(t)}).$$

Choose x' in

$$\dot{f}_{\dot{A}, x_A}(H_{\dot{\alpha}_4, \nu(t)} \cap H_{\dot{\alpha}_5, \nu(t)})$$

but not in $\dot{f}_{\dot{A}, x_A}(\mathbb{A})$. Since u and y both fix x' , so does z . Thus z fixes every point in the interval $[x, x']$. This interval contains points, however, which are in $\dot{f}_{\dot{A}, x_A}(H_{\dot{\alpha}_4, \nu(t)})$ or $\dot{f}_{\dot{A}, x_A}(H_{\dot{\alpha}_5, \nu(t)})$ but not in both. These points are fixed by u or y but not both. Hence they cannot be fixed by z . We conclude that

$$A \cap A^z = \dot{f}_{\dot{A}, x_A}(H_{\alpha, k \cdot \nu(t)}).$$

By 4.8, f_{A, x_A} is the restriction of $\dot{f}_{\dot{A}, x_A}$ to $\mathbb{A}$. Thus $\varphi_\alpha(x_\alpha(0, 0, t)) = k \cdot \nu(t)$ for all $t \in K^*$.

The proof in cases B and F is virtually the same as the proof in case G; we leave the details to the reader. $\square$

By 7.1, there exists a θ -invariant valuation ν_1 such that φ_α satisfies 7.2 or 7.3 with ν_1 in place of ν . By 8.10, it follows that ν_1 is equivalent to ν . Thus 7.11.ii holds. This concludes the proof of 7.11.

* * *

Here, finally, is a precise version of the remark Tits made in [12] that was discussed in the introduction.

Theorem 8.11. *Bruhat-Tits spaces whose building at infinity (in the sense of 4.14 if the rank of Δ is one) is a given Suzuki-Ree building Δ (in the sense of 6.8) defined over a triple (K, θ, L) (in case B) or pair (K, θ) (in cases F or G) are classified by θ -invariant valuations of K . Equivalent θ -invariant valuations of K correspond to equivalent Bruhat-Tits spaces (in the sense of 4.5).*

Proof. Let Δ be a Suzuki-Ree building defined over the triple (K, θ, L) in case B or the pair (K, θ) in cases F or G, and let α and x_α be as in 6.12 (with respect to some apartment Σ of Δ). Suppose that ν is a θ -invariant valuation of K . By 7.4, there exists a valuation φ of the root datum of Δ based at Σ satisfying 7.2 or 7.3. By 3.16, φ is unique up to equipollence. By 4.16, φ determines a unique Bruhat-Tits space $(X, \mathcal{A})$ having Δ as its building at infinity. By 4.15, any valuation equipollent to φ determines the same Bruhat-Tits space.

Suppose, conversely, that $(X, \mathcal{A})$ is a Bruhat-Tits space whose building at infinity is Δ . By 4.16, $(X, \mathcal{A})$ determines a valuation of the root datum of Δ based at Σ which is unique up to equipollence. By 7.1, this equipollence class determines a unique θ -invariant valuation ν of K such that 7.2 or 7.3 holds for some valuation in this equipollence class. $\square$

9. APPENDIX

The inequalities 7.5 and 7.6 hold by 7.4 and the condition (V1). In this section we give an elementary proof of these inequalities which might be of independent interest. In fact, we only give a proof of 7.6; the interested reader will have no trouble applying the same strategy to the inequality 7.5. The proof we give is based on a suggestion of Theo Grundhöfer.

We suppose that we are in case G and that ν is a θ -invariant valuation of K . As in 6.10, we have

$$(9.1) \quad N(r, s, t) = r^{\theta+1}s^\theta - rt^\theta - r^{\theta+3}s - r^2s^2 + s^{\theta+1} + t^2 - r^{2\theta+4}$$

for all (r, s, t) in the group T .

Lemma 9.2. *Let $(r, s, t) \in T$ and suppose that the minimum of $\nu(r)$, $\nu(s)$ and $\nu(t)$ is 0. Then $\nu(N(r, s, t)) = 0$.*

Proof. The Tits endomorphism θ induces a Tits endomorphism of $\bar{K}$. We let $\bar{\theta}$ denote this endomorphism and let $\bar{N}$ be the map obtained by applying the formula 9.1 to the pair $(\bar{K}, \bar{\theta})$ rather than (K, θ) . By 6.11, $\bar{N}$ is anisotropic. $\square$

Lemma 9.3. *Let $(r, s, t) \in T$, let $A = (2\sqrt{3} + 4)\nu(r)$, let $B = (\sqrt{3} + 1)\nu(s)$, let $C = 2\nu(t)$ and let M be the minimum of A , B and C . Then $\nu(N(r, s, t)) = M$.*

Proof. We can assume that $(r, s, t) \neq (0, 0, 0)$. Suppose first that $M = A$. Then

$$N(1, s/r^{\theta+1}, t/r^{\theta+2}) = N(r, s, t)/r^{2\theta+4}$$

by 9.1. Moreover, $\nu(s/r^{\theta+1})$ and $\nu(t/r^{\theta+2})$ are both non-negative since $B \geq A$ and $C \geq A$. Hence

$$\nu(N(1, s/r^{\theta+1}, t/r^{\theta+2})) = 0$$

by 9.2. It then follows that $\nu(N(r, s, t)) = \nu(r^{2\theta+4}) = A = M$.

Suppose next that $M = C$. In this case, we observe that

$$N(r/t^{2-\theta}, s/t^{\theta-1}, 1) = N(r, s, t)/t^2.$$

Moreover, $\nu(r/t^{2-\theta})$ and $\nu(s/t^{\theta-1})$ are both non-negative. Hence

$$\nu(N(r/t^{2-\theta}, s/t^{\theta-1}, 1)) = 0$$

by 9.2. It then follows that $\nu(N(r, s, t)) = \nu(t^2) = C = M$.

It suffices now to assume that $M = B$ and that B is strictly less than both A and C . In this case, $\nu(s^{\theta+1})$ is strictly less than the value under ν of each of the remaining six terms on the right hand side of 9.1. Therefore $\nu(N(r, s, t)) = B = M$. $\square$

Now let $(r, s, t), (w, u, v) \in T$ and let M be the smaller of the two constants obtained by applying 9.3 first to (r, s, t) and then to (w, u, v) . Thus

$$(9.4) \quad \nu(r), \nu(w) \geq M/(2\sqrt{3} + 4)$$

as well as

$$(9.5) \quad \nu(s), \nu(u) \geq M/(\sqrt{3} + 1)$$

and

$$(9.6) \quad \nu(t), \nu(v) \geq M/2.$$

As in 6.10, we have

$$(r, s, t) \cdot (w, u, v) = (r + w, s + u + r^\theta w, t + v - ru + sw - r^{\theta+1}w).$$

Let $a = (2\sqrt{3} + 4)\nu(r + w)$, let $b = (\sqrt{3} + 1)\nu(s + u + r^\theta w)$ and let

$$c = 2\nu(t + v - ru + sw - r^{\theta+1}w).$$

By 9.4–9.6, we have $(2\sqrt{3} + 4)\nu(x) \geq M$ for $x = r$ and $x = w$; $(\sqrt{3} + 1)\nu(x) \geq M$ for $x = s$, $x = u$ and $x = r^\theta w$; and $2\nu(x) \geq M$ for x equal to each of the five terms in the sum

$$t + v - ru + sw - r^{\theta+1}w.$$

Hence $a, b, c \geq M$. By 9.3, therefore,

$$\nu((r, s, t) \cdot (w, u, v)) \geq M = \min\{\nu(N(r, s, t)), \nu(N(w, u, v))\}.$$

REFERENCES

- [1] A. Berenstein and M. Kapovich, Affine buildings for dihedral groups, arXiv:0809.0300.
- [2] F. Bruhat and J. Tits, Groupes réductifs sur un corps local, I. Données radicielles valuées, *Publ. Math. I. H. E. S.* **41** (1972), 5–252.
- [3] T. De Medts, F. Haot, R. Knop and H. Van Maldeghem, On the uniqueness of the unipotent subgroups of some Moufang sets, in *Finite Geometries, Groups, and Computation* (Pingree Park, 2004), pp. 43–66, Walter de Gruyter, Berlin, New York, 2006.
- [4] B. Kleiner and B. Leeb, Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings, *Publ. Math. I. H. E. S.* **86** (1997), 115–197.
- [5] S. Lang, Bruhat-Tits-Räume, *Elem. Math.* **54** (1999), 45–63.
- [6] A. Parreau, Immeubles affines: construction par les normes et étude des isométries, in *Crystallographic Groups and their Generalizations* (Kortrijk, 1999), pp. 263–302, *Contemp. Math.* **262**, Amer. Math. Soc., Providence, 2000.
- [7] M. Ronan, *Lectures on Buildings*, Academic Press, San Diego, 1989.
- [8] G. Rousseau, Euclidean buildings, in *Séminaires et Congrès* **18**, Soc. Math. France, Paris, to appear (<http://hal.archives-ouvertes.fr/hal-00094363>).
- [9] J. Tits, Les groupes simples de Suzuki et de Ree, *Séminaire Bourbaki* **6**, Année 1960–61, pp. 65–82, reprinted by Soc. Math. France, Paris, 1995.
- [10] J. Tits, *Buildings of Spherical Type and Finite BN-Pairs*, Springer Lecture Notes in Math. **386**, 1974.
- [11] J. Tits, Moufang octagons and Ree groups of type 2F_4 , *Amer. J. Math.* **105** (1983), 539–594.
- [12] J. Tits, Immeubles de type affine, in *Buildings and the Geometry of Diagrams* (Como, 1984), pp. 159–190, Lecture Notes in Mathematics **1181**, Springer, 1986.
- [13] J. Tits and R. Weiss, *Moufang Polygons*, Springer Monographs in Math., Springer, Berlin, New York, Heidelberg, 2002.
- [14] R. Weiss, *The Structure of Spherical Buildings*, Princeton University Press, Princeton, 2003.
- [15] R. Weiss, *The Structure of Affine Buildings*, Annals of Math. Studies **168**, Princeton University Press, Princeton, 2009.